

# A Position Paper on Managing Youth Screen Time versus Physical Activity

Reem I. Altamimi, Geoff D. Skinner and Keith V. Nesbitt

*Received 30 Jan 2015 Accepted 26 Feb 2015*

**Abstract**—Childhood obesity is becoming increasingly prolific and problematic. Contributors to childhood obesity include decreased levels of physical activity and increased sedentary behavior. Screen-based entertainment may be an important factor in the development of childhood obesity as children and adolescents prefer to spend time using electronic devices than exercising. While it is difficult to encourage children to cease playing video games, it is possible to change these passive screens to active ones. Information and Communication Technologies (ICTs) have been utilized by academics and researchers to promote levels of physical activity among young people. This paper is part of our continuous research into the use of technology in the facilitation and motivation of children to be more physically active. This paper presents the “MySteps” framework that has been developed to manage youth screen time and exercise performance statistics. By developing technology-based solutions, we intend to increase children and adolescents awareness of their levels of physical activity and screen time. Consequently, self-monitoring and management of screen time and physical activity levels may lead to more active living beginning at an early age and continuing throughout life.

**Keywords**—Active living technologies; eHealth; Monitoring Technologies; Physical Activity

## I. INTRODUCTION

Levels of childhood obesity have been steadily increasing over recent decades [1] [2] [3]. The percentage of obese children in the United States has increased from 7% in 1980 to 18% in 2012 [1]. Although the most recent literature on childhood obesity in the United States shows that there was no significant change in the prevalence of obesity between 2003-2004 and 2011-2012, the overall number of children considered obese remains elevated [4]. Worldwide, the prevalence of overweight and obese children has increased from 4.2% in 1990 to 6.7% in 2010. It is estimated that the prevalence will reach 9.1% by 2020 [2]. These predictions suggest the need for urgent and effective interventions to combat the issue of childhood obesity [2]. Obese children are more likely to become obese adults and therefore children should be specifically targeted, as the benefits of preventing childhood obesity will flow on into adulthood [5] [6] [7]. Obese youth are more likely to have health problems such as cardiovascular disease, diabetes and cancer. Childhood obesity is also associated with social and psychological problems such as depression and low self-esteem [1] [8]. The

cluster of associated medical conditions emphasise the need for the development of effective interventions.

Researchers and health professionals alike are alarmed by the wide spread nature of childhood obesity and its associated health problems and have strived to elucidate causes and contributing factors. Energy intake relative to energy expenditure, genetics, physical activity and sedentary behavior have all be identified as potential causes [3]. Lower levels of physical activity and higher rates of television viewing are two additional important determinants [9]. The integration of technology into our lifestyles and decreased supervision of children by parents who work full time are two factors that may also contribute to decreased levels of physical activity and increased sedentary behavior in children and adolescents [3].

The literature has demonstrated a strong association between time spent in front of screens, especially watching television, and the prevalence of obesity [3] [9] [10]. In one study, with each additional hour of television viewed, the prevalence of obesity increased by 2% [10]. Computers, mobile phones, video game consoles and handheld Internet devices are becoming increasing popular entertaining appliance for children and adolescents. Television stations and video games specifically developed for youth have become a major source of enjoyment. With so much content now available in a variety of media, it is not surprising that children spend a large proportion of their leisure time using electronic entertainment devices [11]. Screen time should be considered an important contributing factor to the increased numbers of obese children [3].

Moreover, researchers have demonstrated an inverse relationship between physical activity level and body fat, particularly in males [12] aged between 6 and 9 years [13]. This may contribute to the development of childhood obesity [13]. Therefore, an important key in the management, prevention and treatment of obesity in children and adolescents may lie in increasing their levels of physical activity [8].

Increasing physical activity and decreasing sedentary behavior has social, emotional and intellectual implications [14]. For example, it reduces anti-social behavior, assists in the development of physical skills, and improves self-esteem and confidence [14]. Regular physical activity is also linked to improved health outcomes and motor skills [15] and reduced risk of many diseases [16] [14]. It contributes to the

---

The primary author is employed and sponsored by Taibah University, Madinah, KSA

prevention of several chronic diseases as the risk of the developing these conditions begins in childhood [16]. Health institutions across the world have set physical activity recommendations for children and adolescents to be at least 60 minutes of moderate-to-vigorous physical activity per day [17] [18] [19]. However, few young people meet this recommendation [19]. Physical activity levels drop between childhood and adolescence and continue to decrease with advancing age [20]. It has been reported that participation in physical activity has declined and sedentary behavior has become more common especially among adolescents [21].

Factors that influence levels of physical activity and may prevent children and adolescents from exercising have been identified. These include, environmental barriers such as limited access to exercise facilities and recreational programs [22], and neighborhood safety [23]. Furthermore, children's motivation is an important determinant of their level of physical activity [24] [22]. Weightless may not be an appropriate motivator for young children to increase their level of physical activity. Young children need to enjoy physical activity in order to engage in it [8]. For example, one mechanism that has been suggested to motivate young children to be more physically active is sport competence and the encouragement of their desire to succeed [24]. Increased participation in physical activity and decreased screen time should be the focus of research into the prevention and treatment of obesity and the promotion of health among children and adolescents [9] [25]. Interventions should promote an active lifestyle and be designed to motivate children to keep fit and active.

Utilising Information and Communication Technologies (ICTs) in physical activity interventions has a positive effect on physical activity behavior change in children and adolescents [26]. ICT-based interventions apply techniques such as Internet or SMS as delivery modes. A major advantage of ICT-based interventions is that they are more readily accessible to a wider audience. Furthermore, these interventions can deliver video, audio and animation content to targeted groups based on their needs and preferences. Other advantages include feedback, automatic reminders and social support [26].

Children and adolescents enjoy using technology and digital devices. A survey of 1,100 parent-child pairs in 2004 showed that the vast majority of US teens used the Internet daily to play games, communicate with friends and seeking information. Some reported using the Internet to search for health, dieting or physical fitness information. Teens are also like to communicate with friends online and share links, photos and videos [27]. Therefore it is expected that developing technology-based solutions for supporting regular participation in physical activity would have positive results.

Our research is designed to explore the best available technology-based solutions for physical activity promotion in children and adolescents. In this paper, we discuss the 'MySteps' solution, which was developed with the aim of managing screen time and physical activity behavior among children and adolescents. This technology-based solution may support the transition of youth behavior away from sedentary lifestyle and towards an active lifestyle. 'MySteps' utilises activity-monitoring technology coupled with web-based feedback. Through this web application, users can

login and monitor their screen time and physical activity performance statistics and compare these to recommended levels. The remainder of the paper provides background information on the current levels of physical activity and sedentary behavior among children and adolescents. The importance of technology-based intervention within this context has also been outlined. The following section details our proposed solution 'MySteps' and its contribution to the active living technology research domain. This paper concludes with our aims and plans to future improve our solution.

## II. SCREEN TIME

### A. Definitions

Sedentary activity refers to activity that involves sitting or lying down [14]. "Screen time" activities are those done in front of a screen, such as watching TV, working on a computer, or playing video games [28]. Screen time is a sedentary activity that requires a very little energy expenditure [28]. Recreational screen time is defined as the time spent in front of screens for entertainment purposes and includes watching TV, playing video games and chatting with friends online [29].

### B. Implications

Spending numerous hours undertaking screen-based activities may contribute to, or increase the risk of, overweight and obesity in children and adolescents [25]. Spending large amounts of time in front of screens reduces the amount of time children spend exercising and contributes to unhealthy eating habits such as snacking [10] [28] [29]. Screen based food advertisements also play a role in encouraging unhealthy eating habits [10] [28] [29]. Furthermore, spending excessive hours in front of screens may lead to sleep disturbance/disorders [30], risk of attention problems, anxiety, and depression [28]. Children and adolescents social skills may also be negatively affected by too many hours spent in front of screens [30].

The association between some specific screen based activities and increased levels of adiposity has been investigated [31]. Reducing television, video and video game use does not significantly affect levels of physical activity [31]. However, it may be a promising method to help prevent childhood obesity as reducing screen time has been shown to affect BMI, triceps skinfold thickness, waist circumference and waist-to-hip ratio [31]. Considering these implications, a screen time limit has been set for children and adolescents. The current guidelines suggest a maximum of two hours a day for children aged between 5 and 18 years [14] [29] [30].

### C. Current Levels

The literature demonstrates a high prevalence sedentary behavior attributable to screen based activities among children and adolescents [25] [32]. Children and adolescents spend an average of 8 hours per day engaged in sedentary behaviors and screen time [25] [33]. This trend is likely to increase as technology advances. Compared to 2003, in 2007, there were fewer children complying with the recommended screen time [34]. There were a number of important findings from a 2010 study on the quantity and nature of media used by young Americans [32]. Firstly, between 2004 and 2009, the amount of time young people

spent using media increased from an average of 6 hours and 21 minutes to 7 hours and 38 minutes per day [32]. Secondly this study revealed that the average time spent using computers, playing video games, watching TV and listening to music and audio increased during this period whereas the use of printed materials such as books declined [32]. Thirdly this study found that the advent of mobile media has facilitates the use of media and expanded the number of hours children and adolescents spend using media. Between 2004 and 2009, the percentage of young people who owned a mobile phone increased from 39% to 66%. In addition, the percentage of young people who own other portable devices such as iPods increased from 18% to 76% in the same time period [32]. Lastly, the number of homes with Internet access expanded from 74% to 84% between 2004 and 2009 [32]. Social networking and video sites like YouTube constitute the majority of the home Internet use. Australian health statistics reveal that in 2011-12; children and adolescents spent greater than two hours per day in front of screens, levels of physical activity decreased and screen-based activity increased as age increased [35]. Additionally, it has been found that children whose parents and caregivers limit media use or set media-related rules spend less time in front of screens [25] [32].

### III. PHYSICAL ACTIVITY

#### A. Definitions

Physical activity refers to any activity that requires physical movement and that results in an increased respiratory and heat rate [14]. Examples include running, playing sports and swimming. Moderate intensity physical activity refers to physical exertion that does not result in breathlessness, for example fast walking or bike riding [14]. Vigorous physical activity refers to activities that require a higher level of physical effort and result in breathlessness, such as running [14].

#### B. Standard Levels of Physical Activity

Children and adolescents are encouraged to participate in 60 minutes of moderate to vigorous physical activity daily [14]. These 60 minutes can be accumulated throughout the day [14]. Rather than measure physical activity in minutes, the criterion-referenced standards for physical activity measure physical activity in steps per day using a pedometer. The median optimal number of steps per day for children aged between 6 and 12 years is 12,000 for girls and 15,000 for boys. It is estimated that this is equivalent to approximately 120 minutes of physical activity per day for girls and 150 minutes per day for boys [36].

#### C. Current Levels of Physical Activity

There have been a number of attempts to assess the level and patterns of physical activity in children and adolescents using objective measurements [37] [20]. However, the lack of a validated tool that captures the multitude of types of physical activity in children is a limitation of research in this area [37]. There are many types of physical activity tracking tools, however, these vary in their mechanisms of measurement and accuracy. There is a need for an accurate device that is able to collect information about children's activity levels. Such a device would assist with assessing whether children meet the recommended levels of physical

activity. This information along with information about what influences and motivates children and adolescents to be physically active would assist with the development of a suitable invention to promote sustained adherence to physically active behavior [37].

The level of physical activity in children and adolescents is a topic of interest among researchers. It has been reported that boys are more physically active than girls [37] [20] [21] [22] and that younger children are more physically active and spend more time engaged in moderate intensity physical activity than adolescences [37] [20]. The 2003-2004 National Health and Nutritional Examination Survey conducted in the United States reported that 42% of children met the recommended level of physical activity compared to 8% of adolescents [20]. Contrastingly, a European study reported that a large majority of young children met the recommended level of physical activity [37]. However, in support of the US study, the European study found that few adolescents met the criteria recommended for physical activity [37]. A study of British children indicated remarkable reductions in physical activity and increases in sedentary, especially among adolescents [21]. Another study reported that only 7% of Canadian children met the daily physical activity recommendations [33]. In a 2011-12 Australian Health Survey of physical activity, only one-third of children, and one in ten young people undertook the recommended 60 minutes of physical activity every day [38]. On average, children and adolescents spend one and a half hours per day physically active [35]. One study that measured physically activity in steps per day using a pedometer reported that children recorded an average of 9,140 Steps/day [35]. Younger children were most likely to achieve the goal of 12,000 steps per day whereas only 7% of children aged 15 to 17 years did so [35].

### IV. A TECHNOLOGICAL SOLUTION

In response to the increased incidence of obesity in children and adolescents, many interventions and treatment programs have been developed with the aim of combatting this problem [39] [15] [40]. These interventions have different focuses including diet and exercise, surgical and drug treatments, behavior change, self-monitoring, and managing screen time and physical activity levels [41] [31] [40]. Several new technologies have been developed and existing technologies exploited for this purpose. For example, Short Message Service (SMS) has been proposed as a method of monitoring children and adolescents' screen time and physical activity behavior [41]. Results showed that children and adolescents seems to be more affected by technology-based programs compared to other traditional paper-based methods of self-monitoring. Using electronic-based programs might be more suitable and provide adherence and acceptability in self-monitoring and behavioral change among young generation [41].

The literature supports the use of Information and Communication Technologies in physical activity interventions as a number of positive effects have been documented for children and adolescents [26]. Electronic tools are useful in increasing individuals' awareness and enhancing their adherence to self-monitoring because of their ability to provide immediate feedback [42]. Another reason

that electronic devices have shown promise in this area is that they have already been embraced by children and adolescents. A survey of a US sample of 1100 parent-child pairs found that 84% of teens owned at least one personal media device: a desktop or laptop computer, mobile phone or Personal Digital Assistant [27]. While personal technological devices are widespread in their use, it remains to be determined which device/s hold the most promise for motivating children and adolescents to adhere to a more physically active lifestyle. It is important that such devices are designed to engage young people, monitor activity accurately and support active lifestyles.

A number of factors that positively affect behavioral change, increase awareness of physical activity and enhance adherence to self-monitoring have been identified, for example home-based tailored interactive programs targeted at children [42] [41]. Additionally, technology that provides immediate feedback and facilitates self-monitoring [41] as well as provides visual presentation of activity level such as charts [42] has been recommended. Studies have demonstrated a strong relationship between parental support and family inclusion, in the form of verbal encouragement or direct assistance, and increased childhood physical activity levels [22] [42] [39].

There are a number of important design considerations in developing technologies that encourage physical activity. Firstly, this technology should give users appropriate credit for their physical activity. There is a problem with currently available activity tracking technologies in that they do not record all types of physical activity. For example, pedometer devices can track steps taken walking or running but cannot monitor activities such as swimming and cycling. Therefore, designers should consider the most common physical activities undertaken by their targeted users [43]. Secondly, technology should allow users to access information regarding their past and current physical activity levels with respect to their goals. Thirdly, technology should include features, such as the ability to share data with friends, which support socializing and communication and which may enhance motivation. Additionally, in developing technology that encourages physical activity, designers should be aware of the practical constraints of users' lifestyles. For example, if technology requires a physical device to be carried or worn, the everyday implications of this should be considered in the design of the device. Finally, the technology should provide reasonable goals that encourage a sustainable increase in physical activity [43].

## V. SIMILAR WORK

A number of similar technology-based projects have been developed with the aim of encouraging and motivating physical activity in specific groups including children, adults [43] and seniors [44]. An example is 'Houston', a prototype mobile phone application that requires its user to wear a pedometer to count their daily steps and then share these step counts with friends. This system has been developed with the aim of encouraging physical activity. Feedback gathered regarding this application reported that it motivated participants to plan increased levels of physical activity [43].

## VI. 'MySteps' ICT FRAMEWORK

### A. Overview of the 'MySteps' Framework

The 'MySteps' solution is part of a larger body of research being conducted to address the ever-growing problem of childhood and adolescent obesity. Our proposed solution focuses on the contributing factors to the issue of increased screen time and decreased physical activity among youth. It is widely accepted that children and adolescents seek entertainment through information and communication technologies. On the basis of an extensive literature review and investigations, we believe that the trend toward an increasing number of overweight and obese children is likely to worsen in the short term. Furthermore, we acknowledge that it is problematic to prevent children playing video games or spending significant time in front of screens. We, therefore, feel that a more useful approach is to utilize the very same technologies they are already prolific in order to encourage a more balanced, active and healthy lifestyle. We aim to employ a technology-based solution to increase awareness of unhealthy behavior and support youth behavior to be more active. Hence, Managing Youth Screen Time and Exercise Performance Statistics (MySteps) provides an easy to access web-based application that integrates a physical exertion tracking device to review an individual's daily activity behavior.

### B. Prototype Design and Implementation

In our web-based platform, we have used the daily step count as a measure of physical activity. We focused on 'recreational screen time' – time that children and adolescents spend in front of screens for fun and entertainment purposes. Our system is composed of two parts: hardware (Fitbit device and a computer) and software ('MySteps' website).

For the hardware component we used a Fitbit tracking device. After extensive product testing and a review of the literature we selected the Fitbit Zip [45]. The Fitbit Zip is a wearable device that can be clipped to pocket, belt, shoes or clothes. The Zip tracks steps taken, calories burned and distance travelled. For our solution we elected to use steps taken. This decision was based on extensive evidence in the literature regarding baseline recommended daily steps for youths (15000 for males and 12000 for females). A computer device, either a desktop or laptop computer with internet access, is necessary to use MySteps. This allows for physical activity and screen time data to be uploaded to the MySteps website.

For the software component, we developed the 'MySteps' website. Users have to register and create an account in order to manage and monitor their screen time and physical activity counts on the 'MySteps' website. Fitbit provides a free API, which we have integrated into the MySteps framework. On registering and logging into MySteps at any given time the user can authenticate with their Fitbit account and their real time exercise statistics are updated in the MySteps application. The screen time element

needs to be self-regulated (parent participation is strongly encouraged for this component of the App) and entered manually directly into the MySteps App. This can be done at any time and any number of times during the day to provide a progressive cumulative total for each day. A history of past behavior is recorded.

Based on screen time and physical activity data, a visual representation is provided. Each element is mapped individually and in combination with the second element against the recommended levels. The graphs are easy to interpret so a young user can view their performance and ideally adjust their behavior for the remainder of the day to meet the recommended levels. Users are also able to monitor their behavior over the preceding week.

## VII. CONCLUSION AND FUTURE WORKS

This paper provides information regarding the increased prevalence of childhood and adolescent obesity and the risk this poses for the future due to the associated health problems. It specifically highlights increasing sedentary behavior and decreasing levels of physical activity among young people as major contributing factors to the development of obesity. Current data relating to levels of physical activity and amount of time children and adolescents spend using screen-based activities is provided. We suggest that technology may help alleviate the problem of obesity among young people. Information and Communication Technologies have been previously utilised for the promotion of physical activity among children and adolescents. We aim to integrate the best available technologies to keep children active. Our previous work in this area found that the use of active video games [46] [47] [48], activity monitoring technologies and a combination of both of these technologies facilitated an active lifestyle among children and adolescents [49] [50] [51] [52]. This paper presents our solution, 'MySteps', a website designed to manage youth screen time and daily step counts.

The MySteps technology framework provides a useful contribution to the encouragement of active living among adolescents and to the active living research domain. Integrating physical activity tracking technology with a web-based application enables youth to track and manage their sedentary and physical activity performance. The solution 'MySteps' offers the benefit of the integration of real-time FitBit activity tracking coupled with self-monitored recreational screen time sedentary time.

The usability of 'MySteps' has been tested by the authors who intend to publish the website and run a small pilot to test its effectiveness in reducing screen time and increasing physical activity among youths. If this pilot demonstrates favourable results, we plan to integrate more features such as a rewards based on the physical exertion. We suggest that the combination of 'MySteps' with exergaming technology would have a further advantage. It is expected that in-game reward and progress based on physical exertion would act as a motivator for greater physical activity in children while simultaneously reducing sedentary time spent in front of screens.

## REFERENCES

- [1] Centers for Disease Control and Prevention, (2014). *Childhood Obesity Facts*. Available: <http://www.cdc.gov/healthyyouth/obesity/facts.htm>
- [2] M. De Onis, M. Blössner, and E. Borghi, "Global prevalence and trends of overweight and obesity among preschool children," *The American journal of clinical nutrition*, vol. 92, pp. 1257-1264, 2010.
- [3] K. F. B. Patricia M. Anderson, "Childhood Obesity: Trends and Potential Causes," *The Future of Children*, vol. 16, pp. 19-45 2006.
- [4] C. L. Ogden, M. D. Carroll, B. K. Kit, and K. M. Flegal, "Prevalence of childhood and adult obesity in the United States, 2011-2012," *JAMA*, vol. 311, pp. 806-814, 2014.
- [5] R. C. Whitaker, M. S. Pepe, J. A. Wright, K. D. Seidel, and W. H. Dietz, "Early adiposity rebound and the risk of adult obesity," *Pediatrics*, vol. 101, pp. e5-e5, 1998.
- [6] W. H. Dietz, "Periods of risk in childhood for the development of adult obesity—what do we need to learn?," *The Journal of nutrition*, vol. 127, pp. 1884S-1886S, 1997.
- [7] R. C. Whitaker, J. A. Wright, M. S. Pepe, K. D. Seidel, and W. H. Dietz, "Predicting obesity in young adulthood from childhood and parental obesity," *New England Journal of Medicine*, vol. 337, pp. 869-873, 1997.
- [8] S. R. Daniels, D. K. Arnett, R. H. Eckel, S. S. Gidding, L. L. Hayman, S. Kumanyika, et al., "Overweight in children and adolescents: pathophysiology, consequences, prevention, and treatment," *Circulation*, vol. 111, pp. 1999-2012, 2005.
- [9] I. Janssen, P. T. Katzmarzyk, W. F. Boyce, C. Vereecken, C. Mulvihill, C. Roberts, et al., "Comparison of overweight and obesity prevalence in school-aged youth from 34 countries and their relationships with physical activity and dietary patterns," *Obesity reviews*, vol. 6, pp. 123-132, 2005.
- [10] W. H. Dietz and S. L. Gortmaker, "Do we fatten our children at the television set? Obesity and television viewing in children and adolescents," *Pediatrics*, vol. 75, pp. 807-812, 1985.
- [11] D. F. Roberts and U. G. Foehr, "Trends in media use," *The future of children*, vol. 18, pp. 11-37, 2008.
- [12] K. Westerterp and M. Goran, "Relationship between physical activity related energy expenditure and body composition: a gender difference," *International journal of obesity and related metabolic disorders: journal of the International Association for the Study of Obesity*, vol. 21, pp. 184-188, 1997.
- [13] E. J. Ball, J. O'Connor, R. Abbott, K. S. Steinbeck, P. S. Davies, C. Wishart, et al., "Total energy expenditure, body fatness, and physical activity in children aged 6-9 y," *The American journal of clinical nutrition*, vol. 74, pp. 524-528, 2001.
- [14] The Department of Health, (June 2014). *Make your move – Sit less – Be active for life!* Available: <http://www.health.gov.au/internet/main/publishing.nsf/content/health-pubhlth-strateg-phys-act-guidelines>
- [15] J. J. Reilly, L. Kelly, C. Montgomery, A. Williamson, A. Fisher, J. H. McColl, et al., "Physical activity to prevent obesity in young children: cluster randomised controlled trial," *Bmj*, vol. 333, p. 1041, 2006.
- [16] D. E. Warburton, C. W. Nicol, and S. S. Bredin, "Health benefits of physical activity: the evidence," *Canadian medical association journal*, vol. 174, pp. 801-809, 2006.

- [17] The Department of Health, (2014). *Australia's Physical Activity and Sedentary Behaviour Guidelines*. Available: <http://www.health.gov.au/internet/main/publishing.nsf/content/health-pubhlth-strateg-phys-act-guidelines>
- [18] W. B. Strong, R. M. Malina, C. J. Blimkie, S. R. Daniels, R. K. Dishman, B. Gutin, *et al.*, "Evidence based physical activity for school-age youth," *The Journal of pediatrics*, vol. 146, pp. 732-737, 2005.
- [19] I. Janssen, "Physical activity guidelines for children and youth This article is part of a supplement entitled Advancing physical activity measurement and guidelines in Canada: a scientific review and evidence-based foundation for the future of Canadian physical activity guidelines co-published by Applied Physiology, Nutrition, and Metabolism and the Canadian Journal of Public Health. It may be cited as Appl. Physiol. Nutr. Metab. 32 (Suppl. 2E) or as Can. J. Public Health 98 (Suppl. 2)," *Applied Physiology, Nutrition, and Metabolism*, vol. 32, pp. S109-121, 2007.
- [20] R. P. Troiano, D. Berrigan, K. W. Dodd, L. C. Masse, T. Tilert, and M. McDowell, "Physical activity in the United States measured by accelerometer," *Medicine and science in sports and exercise*, vol. 40, p. 181, 2008.
- [21] N. H. Brodersen, A. Steptoe, D. R. Boniface, and J. Wardle, "Trends in physical activity and sedentary behaviour in adolescence: ethnic and socioeconomic differences," *British journal of sports medicine*, vol. 41, pp. 140-144, 2007.
- [22] J. F. Sallis, J. J. Prochaska, and W. C. Taylor, "A review of correlates of physical activity of children and adolescents," *Medicine and science in sports and exercise*, vol. 32, pp. 963-975, 2000.
- [23] B. E. Molnar, S. L. Gortmaker, F. C. Bull, and S. L. Buka, "Unsafe to play? Neighborhood disorder and lack of safety predict reduced physical activity among urban children and adolescents," *American Journal of Health Promotion*, vol. 18, pp. 378-386, 2004.
- [24] S. Biddle and M. Goudas, "Analysis of children's physical activity and its association with adult encouragement and social cognitive variables," *Journal of School Health*, vol. 66, pp. 75-78, 1996.
- [25] D. W. Lou, "Sedentary Behaviors and Youth: Current Trends and the Impact on Health," ed: Active Living Research, January 2014.
- [26] P. W. Lau, E. Y. Lau, D. P. Wong, and L. Ransdell, "A systematic review of information and communication technology-based interventions for promoting physical activity behavior change in children and adolescents," *Journal of medical Internet research*, vol. 13, 2011.
- [27] A. Lenhart, M. Madden, and P. Hitlin, "Teens and technology: Youth are leading the transition to a fully wired and mobile nation," 2005.
- [28] N. K. Kaneshiro, (2013). *Screen time and children*. Available: <http://www.nlm.nih.gov/medlineplus/ency/patientinstructions/000355.htm>
- [29] The Parent's Jury, (2013). *Recreational Screen Time* Available: <https://www.parentsjury.org.au/key-concerns/physical-activity/recreational-screen-time>
- [30] Healthy Kids, (2015). *Switch off the Screen*. Available: <http://www.healthykids.nsw.gov.au/kids-teens/switch-off-the-screen.aspx>
- [31] T. N. Robinson, "Reducing children's television viewing to prevent obesity: a randomized controlled trial," *Jama*, vol. 282, pp. 1561-1567, 1999.
- [32] V. J. Rideout, U. G. Foehr, and D. F. Roberts, "Generation M [superscript 2]: Media in the Lives of 8-to 18-Year-Olds," *Henry J. Kaiser Family Foundation*, 2010.
- [33] R. Wilson, "Promoting physical activity," *BCMJJ*, vol. 54, p. 335, September 2012 2012.
- [34] S. B. Sisson, S. T. Broyles, D. R. Brittain, and K. Short, "Obesogenic behaviors in US school children across geographic regions from 2003-2007," *Open Journal of Preventive Medicine*, vol. 1, p. 25, 2011.
- [35] Australian Bureau of Statistics, (2013). *Australian Health Survey: Physical Activity, 2011-12* Available: <http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/4364.0.55.004Chapter1002011-12>
- [36] C. Tudor-Locke, R. P. Pangrazi, C. B. Corbin, W. J. Rutherford, S. D. Vincent, A. Raustorp, *et al.*, "BMI-referenced standards for recommended pedometer-determined steps/day in children," *Preventive medicine*, vol. 38, pp. 857-864, 2004.
- [37] C. J. Riddoch, L. B. Andersen, N. Wedderkopp, M. Harro, L. Klasson-Heggebo, L. B. Sardinha, *et al.*, "Physical activity levels and patterns of 9-and 15-yr-old European children," *Medicine and science in sports and exercise*, vol. 36, pp. 86-92, 2004.
- [38] The Department of Health, (2014). *Research and Statistics*. Available: <http://www.health.gov.au/internet/main/publishing.nsf/Content/health-pubhlth-strateg-active-evidence.htm>
- [39] L. H. Epstein, A. Valoski, R. R. Wing, and J. McCurley, "Ten-year outcomes of behavioral family-based treatment for childhood obesity," *Health Psychol*, vol. 13, pp. 373-83, Sep 1994.
- [40] L. H. Epstein, M. D. Myers, H. A. Raynor, and B. E. Saelens, "Treatment of pediatric obesity," *Pediatrics*, vol. 101, pp. 554-570, 1998.
- [41] J. R. Shapiro, S. Bauer, R. M. Hamer, H. Kordy, D. Ward, and C. M. Bulik, "Use of text messaging for monitoring sugar-sweetened beverages, physical activity, and screen time in children: a pilot study," *Journal of nutrition education and behavior*, vol. 40, pp. 385-391, 2008.
- [42] S. J. Bartlett, P. Lukk, A. Butz, F. Lampros-Klein, and C. S. Rand, "Enhancing medication adherence among inner-city children with asthma: results from pilot studies," *Journal of Asthma*, vol. 39, pp. 47-54, 2002.
- [43] S. Consolvo, K. Everitt, I. Smith, and J. A. Landay, "Design requirements for technologies that encourage physical activity," in *Proceedings of the SIGCHI conference on Human Factors in computing systems*, 2006, pp. 457-466.
- [44] T. W. Bickmore, L. Caruso, and K. Clough-Gorr, "Acceptance and usability of a relational agent interface by urban older adults," in *CHI'05 extended abstracts on Human factors in computing systems*, 2005, pp. 1212-1215.
- [45] Fitbit. (2015). Available: <http://www.fitbit.com/au/home>
- [46] R. Altamimi and G. Skinner, "A Survey of Active Video Game Literature," *Journal of Computer and Information Technology*, vol. 1, pp. 20-35, 2012.
- [47] P. Ilung, R. Altamimi, and G. D. Skinner, "Exertainment: Designing Active Video Games to Get Youth Moving," in *Proceedings of the International MultiConference of Engineers and Computer Scientists*, 2014.
- [48] R. A. Ilung Pranata, Geoff Skinner, "An Adaptive Framework Allowing Active Video Games to Address Child Obesity," presented at the 7th Annual International Conference on Computer Games, Multimedia and Allied Technology 2014, Singapore, 2014.

- [49] R. Altamimi, G. Skinner, and K. Nesbitt, "FITTER-A Framework for Integrating Activity Tracking Technologies into Electric Recreation for Children and Adolescents."
- [50] R. Altamimi, G. Skinner, and K. Nesbitt, "A Focused Review and Initial Conceptual Design for Merging Exergame and Activity Monitoring Technologies," in *Entertainment Computing-ICEC 2014*, ed: Springer, 2014, pp. 77-83.
- [51] G. S. Reem Altamimi, Keith Nesbitt, "Pitfalls and Promises of Exergaming and Activity Monitoring Technologies," presented at the The 9th International Conference on Information Technology and Applications (ICITA2014), Sydney, Australia, 2014.
- [52] G. Skinner and R. Altamimi, "Active Video Games (AVG) For Promoting E-Health Initiatives-Integration Of Contemporary Information And Communication Technologies (ICT)."

#### AUTHOR'S PROFILE

**Reem I. Altamimi** is a PhD Candidate at the University of Newcastle, NSW, Australia. She is employed and sponsored by Taibah University, Madinah, Saudi Arabia. The author earned her Bachelor degree in Computer Science from Taibah University, Madinah, Saudi Arabia, in 2006. She obtained a Master of Information Technology (Advanced) with Honours Class I from the University of Newcastle, Australia, graduating in 2012. The Author is particularly interested in Active Living Technologies. Her research is based on exploring the integration of the best available contemporary technologies in order to facilitate and motivate active living among children and adolescents.

**Dr Geoff Skinner** is Head of the IT Discipline at the University of Newcastle, and member of the Priority Research Centre for Physical Activity and Nutrition. He has close to 100 International peer reviewed publications and 2 book chapters in the field of Data Security. Coupled with over 15 years professional Industry experience, he brings a wealth of knowledge to his chosen research domains. These include Active Living Technologies, Exertainment, Data and Information Security, and eLearning.

**Dr Keith Nesbitt** is a Senior Lecturer in Information Technology at the University of Newcastle. His research interests include computer games, virtual environments, user-interface design, information visualisation, agent-based modelling and creative processes. With an applied focus in these areas of research, Dr Nesbitt explores how these elements can be employed to solve real world problems. In particular, he looks at how information can be shared and received through engaging the senses. He has successfully secured funding for various projects, including the Strategic Initiatives Research Fund grant for his work in health informatics; the Teaching and Learning Project grant for the development of a virtual pharmacy; and the PRC in Physical Activity and Nutrition Seed Grant for his research in quantifying dysfunctional movement. Dr Nesbitt previously worked as a research scientist for BHP, assessing emerging computing technology and their potential impact on BHP business domains. He currently teaches into the courses related to computer games and computer programming. Dr Nesbitt is also a member of the School of Design, Communication and Information Technology's research committee.

***This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.***