



NOVA

University of Newcastle Research Online

nova.newcastle.edu.au

Costigan, Sarah A.; Lubans, David R.; Lonsdale, Chris; Sanders, Taren & del Pozo Cruz, Borja. "Associations between physical activity intensity and well-being in adolescents" Published in *Preventive Medicine*, Vol. 125, p. 55-61, (2019).

Available from: <http://dx.doi.org/10.1016/j.ypmed.2019.05.009>

© 2019. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Accessed from: <http://hdl.handle.net/1959.13/1403214>

Associations between physical activity intensity and well-being in adolescents

Sarah A. Costigan^{1,2}, David R. Lubans¹, Chris Lonsdale³, Taren Sanders³ and Borja del Pozo Cruz³

¹ Priority Research Centre in Physical Activity and Nutrition, University of Newcastle, Newcastle, NSW, Australia

² Deakin University, Geelong, School of Exercise and Nutrition Sciences, Australia

³ Institute for Positive Psychology, Faculty of Health Sciences, Australian Catholic University, North Sydney, NSW, Australia

Corresponding Author

Sarah Costigan

Email: Sarah.Costigan@newcastle.edu.au

Address: Priority Research Centre in Physical Activity and Nutrition, The University of Newcastle, University Drive, Callaghan NSW 2308

Phone: +61 3 522 73461

Word count

Abstract: 226

Main text: 3433

ABSTRACT

This study aims to explore associations between physical activity intensity and well-being (i.e., positive and negative affect) in adolescents. A secondary aim was to determine if associations were moderated by sex. Grade 8 students from 14 government-funded secondary schools in low socio-economic areas of Western Sydney (Australia) were assessed. Data from three timepoints (baseline, 7-8 months, and 14-15 months) were combined to increase the sample size. Physical activity was objectively assessed for 1-week at each timepoint using Actigraph accelerometers. Time (minutes/day) in light, moderate and vigorous physical activity was estimated. The short form Positive and Negative Affect Scale for Children was used to measure well-being. Quantile regression was used to analyse the data. A total of 3,140 observations were collected from 1,223 students (mean age at baseline: 12.9(0.54); 55.1% male). Light and moderate physical activity were not associated with well-being. Higher levels of vigorous physical activity were associated with more positive affect [β (SE)=0.307 (0.06), $p<0.001$], to an estimated vigorous physical activity turning point [Point(95%CI)=36.48 mins/day (31.39-41.59)]. Similarly, higher levels of vigorous physical activity were associated with less negative affect [β (SE)=-0.250 (0.06), $p<0.001$] up to the estimated vigorous physical activity turning point [Point(95%CI)=37.35 mins/day (31.27-43.44)]. The negative association between vigorous physical activity and negative affect was more pronounced in females than in males. Our findings highlight the importance of adolescents engaging in vigorous physical activity to improve positive affect and reduce negative affect.

Keywords: adolescents, vigorous physical activity, well-being, affect

INTRODUCTION

The physical, mental and cognitive health benefits of engaging in physical activity for young people are well documented (Janssen & LeBlanc, 2010). To gain these benefits, international physical activity guidelines recommend children (5-12 years) and adolescents (13–17 years) need to accrue at least 60 minutes of moderate-to-vigorous physical activity (MVPA) daily (World Health Organization, 2012). However, fewer than 20% of adolescents meet these targets (Rhodes, Janssen, Bredin, Warburton, & Bauman, 2017). In 2012, the World Health Organization (WHO) Physical Activity Guidelines were revised to include recommendations for youth to engage in vigorous physical activity on three days each week (World Health Organization, 2012).

Emerging evidence indicates that vigorous intensity physical activity is more strongly associated with physical health outcomes in young people, compared to physical activity of light and moderate intensity (Gutin & Owens, 2011). For instance, a 7-year longitudinal study found vigorous physical activity (but not MVPA) to be inversely associated with BMI in a nationally representative sample of young people ($n=4,770$; aged 7 at baseline) (Hamer & Stamatakis, 2018). Furthermore, a cross-sectional study ($n=605$; mean age 12.1 years) of objectively measured physical activity intensity and cardio-metabolic risk factors reported vigorous physical activity (but not light or moderate) was associated with reduced waist circumference, BMI-Z, systolic blood pressure, and improved cardiorespiratory fitness (Hay et al., 2012). While evidence suggests vigorous physical activity is important for physical health, less is known regarding the effect on mental health outcomes.

Mental health is not merely the absence of disease, the WHO broadly defines mental health as *“a state of well-being in which the individual realizes his or her own abilities, can cope with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to his or her community”* (World Health Organization, 2004). It has been proposed that mental health can only be sufficiently represented when considering the

interaction of both negative (e.g., ill-being, anxiety, depression, perceived stress) and positive (e.g., well-being, self-esteem, resilience) indicators (Masse et al., 1998). Of note, there is conceptual overlap among the different negative and positive indicators of mental health (Ryan & Deci, 2001). For the purpose of this paper, the term well-being will be used to reflect students' self-reported affect.

Review-level evidence suggests physical activity has the potential to improve aspects of adolescents' well-being. A review of reviews reported small-to-moderate positive effects of physical activity on self-esteem (Biddle & Asare, 2011), and a review by Lubans and colleagues (2012) found evidence to suggest that outdoor adventure programs can improve self-concept, resilience, perceptions of alienation and self-control (Lubans, Plotnikoff, & Lubans, 2012). However, much of the available evidence is derived from low quality studies and evidence relies heavily on self-reported physical activity. The few studies using objective measures of physical activity have typically involved small samples (Poitras et al., 2016), furthermore methodological challenges associated with use of different accelerometer cutpoints may impact consistency of findings (Cain, Sallis, Conway, Van Dyck, & Calhoun, 2013). Most studies have explored linear relationships, however studies in other population groups suggest that this relationship may not be linear, and therefore other statistical methods such as quantile regression may be appropriate. For example, quantile regression has been used to examine the relationship between physical activity and obesity in youth aged 6-18 years (Mitchell et al., 2017) and the association between physical activity and teacher-reported academic performance in 9-10 year olds (Zhang et al., 2015). In addition to these methodological shortcomings, there is an absence of research specifically examining the relationship between physical activity intensity and affective well-being amongst adolescents (Parfitt, Pavey, & Rowlands, 2009).

To address these research gaps, our study aims to examine associations between physical activity intensity and hedonic well-being (i.e., positive and negative affect) in adolescents using quantile regression, exploring non-linear relationships. A secondary aim was to determine if associations are moderated by sex, as previous research has consistently documented that females engage in less physical activity than males (Belcher et al., 2010; Lagerberg, 2005), and suggests males and females may experience differential mental health effects (Kremers, de Bruijn, Droomers, van Lenthe, & Brug, 2007). Based on previous research (Biddle & Asare, 2011), we hypothesised that: (i) vigorous physical activity will be associated with well-being, (ii) light and moderate physical activity will not be associated with well-being, and (iii) the association between physical activity intensity and well-being will be moderated by sex.

METHODS

Data from the ‘Activity and Motivation in Physical Education (AMPED)’ cluster RCT were examined in the current study. The full AMPED methodology is published elsewhere (Lonsdale et al., 2016). Briefly, students in Grade 8 from 14 government-funded secondary schools in Western Sydney (Australia) were recruited to participate in an intervention aiming to increase opportunities for physical activity in physical education (PE) and improve motivation for physical activity and PE. Data from three timepoints (baseline, 7-8 months, and 14-15 months follow-up) were combined to increase the sample size. Ethical approval were obtained from the Australian Catholic University (2014185N) and the New South Wales Department of Education (2013162).

Measures

Participants self-reported birthdate, sex, country of birth, language spoken at home and Indigenous origin. Socioeconomic status was measured using the Family Affluence Scale

(Currie et al., 2008). Research assistants measured weight (recorded to the nearest 0.1 kg) and height (recorded to the nearest 0.1 cm), and body mass index (BMI) was calculated ($\text{weight(kg)}/\text{height (m)}^2$). BMI-z scores were calculated using the LMS (lambda-mu-sigma) method (Flegal & Cole, 2013).

Physical activity

Physical activity was assessed using ActiGraph accelerometers (GT1M, GT3X, GT3X+ models; Fort Walton Beach, FL). A range of device models were included as only vertical axis output were used for the current study, and other studies in adolescent populations have reported similar vertical axis outputs between devices (John, Sasaki, & Freedson, 2010). Study participants wore accelerometers for one week (five weekdays, two weekend days) at each timepoint (baseline, 7-8 months, 14-15 months), accelerometers were initialised to record data at 1 second epochs (Lonsdale et al., 2015). Consistent with other studies in adolescent populations (Cain et al., 2013), to be included in the analyses valid data defined as wear time of >8 hours on 3 days or more (including 2 weekdays) was required. Time (minutes/day) spent in physical activity of differing intensities (light (101–2295 counts/minute), moderate (2296–4011 counts/minute) and vigorous (>4012 counts/minute)) was estimated, using validated cut-points (Evenson, Catellier, Gill, Ondrak, & McMurray, 2008). Non-wear time, defined as ≥ 30 minutes of continuous '0' counts, was removed from the dataset (Carson, Cliff, Janssen, & Okely, 2013).

Well-being

Well-being was assessed at each timepoint (baseline, 7-8 months, and 14-15 months follow-up) using the Positive and Negative Affect Scale (PANAS) for Children (Ebesutani et al., 2012). The PANAS is a self-report questionnaire using a 10-item scales to measure positive and negative affect. Using a Likert scale of 1 (*not at all*) to 5 (*very much*), respondents rated

each adjective according to the extent to which each describes the way they have felt during the ‘past few weeks’ (*positive affect*: joyful, cheerful, happy, lively, proud; *negative affect*: miserable, mad, afraid, scared, sad). Scores of the five-items were summed to create a total score out of 25 for each scale. Scores ranged from 5–25, higher positive affect scale scores representing higher positive affect, and lower negative affect scale scores representing reduced negative affect. Studies examining youth have found the PANAS to be a reliable and valid instrument to assess affect (Laurent et al., 1999).

Data analysis

Statistical analyses were conducted using R software (R project version 3.5.1). Summary statistics described variables of interest. Significance levels were set at $p < 0.05$. Participants who provided physical activity and affect data from at least one timepoint were included in the analysis. Polynomial (quadratic) mixed models were used to capture the potentially non-linear relationships between physical activity of light, moderate and vigorous intensity and positive and negative affect. An individual-level random term was also added to the models in order to account for the potential individual variability. When a quadratic effect was detected, a series of sensitivity checks were performed. The turning point of each curve and its 95% confidence interval (CI) was calculated using the Delta Method (Francq & Cartiaux, 2016). This approach has been used before in similar studies (Owen, Parker, Astell-Burt, & Lonsdale, 2018). Potential moderation effects (sex) were tested by including an interaction term in the equations. Lastly, quantile regressions were used in order to capture the potential differences in the impact that physical activity can have across the positive and negative affect distribution (we tested the orthogonal polynomial relationship at the 25th, 50th, 75th and 90th percentiles) in the study sample. All models were adjusted for sex, BMI-z score, ethnicity, wear-time, and other physical activity intensities (light, moderate,

vigorous). Standardised regression coefficients of 0.1, 0.3 and 0.5 were considered small, moderate and large, respectively (Cohen, 1992).

RESULTS

Baseline participant characteristics

Briefly, the study sample included 1,223 adolescents providing 3,140 observations across three timepoints. At baseline the sample was 55.1% male and mean age of 12.94(0.54) years. The sample included 50% classified as ‘normal weight’ (mean BMI: 22.15(4.89)). Mean accelerometer wear time across the three timepoints was 709.94(120.38) minutes per day, including 81.51(27.21), 27.95(11.07) and 18.56(10.91) minutes of daily light, moderate and vigorous physical activity, respectively. Mean scores of 18.05(4.52) and 9.33(4.09) were recorded for positive and negative affect, respectively.

Positive affect and physical activity

There were no linear or quadratic relationships detected between light or moderate intensity physical activity and positive affect. Figure 1 shows the curvilinear relationship between vigorous physical activity and positive affect in the study sample. Increasing levels of vigorous physical activity had a positive and moderate association with positive affect [β (SE)=0.307 (0.06), $p<0.001$] to the estimated vigorous physical activity turning point [Point(95%CI)=36.48min/day (31.39-41.59)], after which positive affect levels were lower with increasing levels of vigorous physical activity [β (SE)=-0.210 (0.053), $p<0.001$]. Of the 3,140 observations, 8.56% included a vigorous physical activity volume above this turning point. Sensitivity analyses showed sex had no effect on this curvilinear relationship (Figure 2). The curvilinear relationship between vigorous physical activity and positive affect was consistent across the entire spectrum of positive affect levels (Figure 3).

Negative affect and physical activity

There were no linear or quadratic relationships detected between light or moderate intensity physical activity and negative affect. As shown in Figure 1, a curvilinear relationship existed, vigorous physical activity had a moderate and negative association with negative affect [$\beta(\text{SE})=-0.3 (0.06)$, $p<0.001$] up to the estimated vigorous physical activity turning point [Point(95%CI)=37.35mins/day (31.27-43.44)], after which negative affect levels were higher [$\beta(\text{SE})=0.184(0.05)$, $p=0.002$]. Of the 3,140 observations, 7.73% included vigorous physical activity volume above this turning point.

Sex sensitivity analysis showed for the decreasing part of the curve ($p=0.03$) the association between vigorous physical activity and negative affect was more pronounced in females than males ($\beta(\text{SE})=-0.038(0.02)$); however, no sex effect was detected for the increasing part of the curve (Figure 2). This means that females appeared to benefit more from vigorous physical activity up to the turning point, but showed similar negative affect as males when engaging in very high levels of vigorous physical activity. Quantile regression analysis suggested that the curvilinear relationship between vigorous physical activity and negative affect was statistically significant for the 25th, 50th, and 75th percentiles but not for the 90th percentile of negative affect (Figure 3), and that regression coefficients for the quantile 90th was statistically significant different from the other quantiles ($p<0.05$).

DISCUSSION

This study examined associations between physical activity intensity and well-being (positive and negative affect) in a large sample of Australian adolescents. Light and moderate physical activity were not associated with well-being. A favourable moderate association was observed between vigorous physical activity and more positive affect, to a turning point of ~36 minutes after which less positive affect was observed. When examining negative affect, engaging in up to ~37 minutes of vigorous physical activity was associated with less negative affect. The association between vigorous physical activity and negative affect was more

pronounced in females than in males. Our study findings provide evidence for the benefits of engaging in vigorous physical activity.

Positive affect

In the current study, no associations between light or moderate intensity physical activity and affect were evident, however a curvilinear relationship between vigorous physical activity and positive affect was observed. Engaging in up to ~36 minutes/day of vigorous physical activity was associated with more positive affect. To our knowledge, there is an abundance of reviews examining associations between physical activity and various mental health outcomes (Biddle & Asare, 2011; Spruit, Assink, van Vugt, van der Put, & Stams, 2016; Wu et al., 2017), however there does not currently appear to be a review or meta-analysis specifically investigating the effect of physical activity on well-being outcomes in adolescents. While available evidence suggests physical activity to be beneficial for adolescent's self-esteem (Biddle & Asare, 2011), self-concept (Lubans et al., 2012; Spruit et al., 2016), externalizing problems and internalizing problems (Spruit et al., 2016), and health-related quality of life (Wu et al., 2017), the effect of physical activity intensity has scarcely been reviewed. Of note, emerging evidence suggests vigorous physical activity in the form of high intensity interval training (HIIT) has both short- and long-term benefits for adolescents' mental health, specifically after an 8-week HIIT trial small intervention effects were observed for well-being outcomes (Costigan, Eather, Plotnikoff, Hillman, & Lubans, 2016).

Negative affect

In the current study, high levels of vigorous physical activity were associated with less negative affect, until the turning point of ~37 minutes after which more negative affect was observed. To date, most studies investigating associations between physical activity and well-being have been cross-sectional, and have reported mixed findings. For example, Goldfield

and colleagues (2011) reported a significant reduction in depressive symptoms in males and anxiety for females ($n=1259$; age: 14.8 ± 1.8) who self-reported engaging in greater amounts of vigorous physical activity (defined as number of times/week engaging in vigorous physical activity for more than 15 minutes/day) (Goldfield et al., 2011). However, other cross-sectional studies have found no association between vigorous physical activity and depression or anxiety (Allison et al., 2005), likewise in baseline data from the Trial of Activity for Adolescent Girls ($n=1,397$ girls; 12 age) no significant associations between depressive symptoms and physical activity intensity were observed (Johnson et al., 2008). Mixed findings may be attributed to differences in physical activity measurement for example self-report, small sample sizes in some studies and analysis approached utilised (e.g., the majority of studies have explored linear relationships, however relationships may not be linear across the distribution of the x variable).

Vigorous physical activity turning point

In the current study, a curvilinear relationship between time spent in vigorous physical activity and affect was observed. Interestingly, in a small number of cases (~7% for positive affect and ~8% for negative affect), engaging in higher amounts of vigorous activity was associated with less positive affect (i.e., >36 minutes/day) and more negative affect (i.e., >37 minutes/day). While engaging in regular physical activity has been shown as beneficial for adolescent health and well-being (Poitras et al., 2016), the ‘extreme exercise hypothesis’ (Eijssvogels, Thompson, & Franklin, 2018) suggests when performed excessively or for extended periods (i.e., beyond optimal dose), exercise can adversely effect health. Previous studies indicate that those engaging in excessive levels of physical activity (e.g., elite athletes) can suffer from a range of other factors, including over-training, injury, stress, and pressure to perform (Armstrong & Vanheest, 2002), which may contribute to poor physical (Eijssvogels et al., 2018) and mental health outcomes. Furthermore, the curvilinear relationship

between vigorous physical activity and negative affect was statistically significant for the 25th, 50th, and 75th percentiles, however not for the 90th percentile. It is likely that multiple factors contribute to these findings, for instance those reporting high negative affect may be experiencing other co-occurring conditions, adolescents participating in greatest vigorous physical activity (e.g., elite athletes) were experiencing poor mental health, or the environment, activities and/or social conditions may not be conducive to lowering negative affect (Rhodes & Kates, 2015). Longitudinal evidence is needed to assess the temporal sequence.

Potential mechanisms responsible for the effect of physical activity on mental health

A range of neurobiological, psychosocial and behavioural mechanisms might explain the relationship between intensity and well-being. Intense physical activity is likely to cause a release of endorphins which may result in changes to affect. Indeed, the ‘endorphin hypothesis’ proposes a reduction in depression and anxiety due to endogenous opioid peptide secretions released in the brain (Dishman & O'Connor, 2009). Adult studies have highlighted the potential role of endocannabinoids in producing exercise-induced improvements in affect, well-being, anxiety, post-exercise calm and pain sensation (Dietrich & McDaniel, 2004). Interestingly, the neurobiological effects of exercise appear to exhibit a ‘u-shaped’ curve, whereby moderate intensity physical activity resulted in greatest mental health effects, while highest and lowest intensity physical activity yielded limited effects (Raichlen, Foster, Seillier, Giuffrida, & Gerdeman, 2013).

A number of psychosocial mechanisms may also explain the positive association of physical activity on well-being, including social interaction, mastery in the physical domain, and improvements to appearance/self-perceptions (Morgan, Young, Smith, & Lubans, 2016). While there is a lack of research examining such underlying mechanisms for adolescent

populations, participation in physical activity programs that involve bouts of vigorous physical activity have the potential to improve adolescents' well-being via changes in fitness and body shape, improved self-efficacy and physical self-concept. Notably, a review of the psychological and social benefits of sport participation found adolescents participating in sport scored higher for psychological domains of well-being (Eime, Young, Harvey, Charity, & Payne, 2013). The 'social interaction hypothesis' posits that interactions and mutual support gained via participating in physical activity together (North, McCullagh, & Tran, 1990), and may explain the beneficial effects of vigorous physical activity on well-being. It has been hypothesised that changes in well-being associated with physical activity may be mediated by behavioural mechanisms such as improved sleep, coping skills and self-regulation (Morgan et al., 2016).

Moderators

Sex was a moderator of the association between vigorous physical activity and negative affect. Females appeared to benefit most from engaging in vigorous intensity physical activity (i.e., higher vigorous physical activity was associated with less negative affect) up to the turning point. Previous research suggests sex may be a key factor moderating associations between physical activity and well-being (Kremers et al., 2007). A more pronounced reduction in negative affect for females might be explained by existing higher negative affect scores compared to males (PANAS negative affect baseline score: 10 vs. 9.2, $p < 0.05$) and lower levels of vigorous physical activity reported by females compared to males (14.7 mins/day vs. 22.03 min/day, $p < 0.05$). Sex differences in physical activity levels are consistently documented, with females engaging in less physical activity than males (Belcher et al., 2010; Lagerberg, 2005), this trend continues when specifically examining vigorous physical activity participation (Parfitt et al., 2009). Alternatively, physical activity may yield other improvements for females (e.g., self-esteem, self-concept and appearance), which also

contribute to less negative affect and more positive affect. Strategies tailored specifically according to sex, which promote relevant opportunities (e.g., activity type, delivery methods) to engage in more vigorous physical activity, appear needed for inactive girls. The importance of targeted and tailored physical activity interventions for enhancing well-being, which reflect sex differences, has been previously highlighted (Morgan et al., 2016).

Strengths and limitations

This study addresses gaps highlighted in existing literature and has a number of strengths. Firstly, the use of objectively measured physical activity using accelerometry is a strength, as much of the existing literature is based on self-reported physical activity. Secondly, this study examined associations between affect and time spent in physical activity of differing intensities (light, moderate, vigorous). Finally, the large number of observations included in the analyses is a strength, previous literature has commonly included small samples. However, study limitations should also be acknowledged. The cross-sectional design of this study limits causal inferences, future experimental studies are encouraged to further investigate associations. The sample is not representative of the wider population which limits generalizability of the study findings. Furthermore, affect is just one component of well-being and a range of components, including indicators of eudemonic wellbeing (e.g., life satisfaction) should be tested in the future to provide a more robust indication of changes in this aspect of mental health.

Conclusions and future direction

The evidence from this study adds to the current body of knowledge documenting the potential benefits of vigorous physical activity for adolescents' well-being. More time spent in vigorous physical activity was associated with improved affective well-being. However, for a small number of adolescents extended vigorous intensity physical activity may heighten

negative affect and lessen positive affect. Females appear to benefit most from engaging in vigorous physical activity, which may be explained by lower vigorous physical activity levels and higher negative affect reported. While participation in any physical activity should be supported, the current study highlights the importance of encouraging adolescents to participate in bouts of vigorous intensity physical activity for enhanced affective wellbeing. Our findings help to provide evidence for necessary vigorous physical activity accrual for adolescents, as recommended on 3 days per week in the current physical activity guidelines.

Acknowledgements

An Australian Research Council Discovery Grant (DP130104659) funded this research.

DRL is supported by an Australian Research Council (FT140100399) Future Fellowship.

The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation, and do not constitute endorsement by ACSM.

Conflict of Interest

No potential conflict of interest was reported by the authors.

References

- Allison, K. R., Adlaf, E. M., Irving, H. M., Hatch, J. L., Smith, T. F., Dwyer, J. J., & Goodman, J. (2005). Relationship of vigorous physical activity to psychologic distress among adolescents. *J Adolesc Health, 37*(2), 164-166.
- Armstrong, L. E., & Vanheest, J. L. (2002). The unknown mechanism of the overtraining syndrome. *Sports Med, 32*(3), 185-209.
- Belcher, B. R., Berrigan, D., Dodd, K. W., Emken, B. A., Chou, C.-P., & Spuijt-Metz, D. (2010). Physical activity in US youth: Impact of race/ethnicity, age, gender, & weight status. *MSSE, 42*(12), 2211.
- Biddle, S. J., & Asare, M. (2011). Physical activity and mental health in children and adolescents: a review of reviews. *BJSM, bjsports90185*.
- Cain, K. L., Sallis, J. F., Conway, T. L., Van Dyck, D., & Calhoun, L. (2013). Using accelerometers in youth physical activity studies: a review of methods. *Journal of Physical Activity and Health, 10*(3), 437-450.
- Carson, V., Cliff, D. P., Janssen, X., & Okely, A. D. (2013). Longitudinal levels and bouts of sedentary time among adolescent girls. *BMC pediatrics, 13*(1), 173.
- Cohen, J. (1992). A power primer. *Psych bull, 112*(1), 155.
- Costigan, S. A., Eather, N., Plotnikoff, R. C., Hillman, C. H., & Lubans, D. R. (2016). High-intensity interval training for cognitive and mental health in adolescents. *MSSE, 48*(10), 1985-1993.
- Currie, C., Molcho, M., Boyce, W., Holstein, B., Torsheim, T., & Richter, M. (2008). Researching health inequalities in adolescents: the development of the Health Behaviour in School-Aged Children (HBSC) family affluence scale. *Social science & medicine, 66*(6), 1429-1436.
- Dietrich, A., & McDaniel, W. F. (2004). Endocannabinoids and exercise. *BJSM, 38*(5), 536-541.
- Dishman, R. K., & O'Connor, P. J. (2009). Lessons in exercise neurobiology: the case of endorphins. *MHPH, 2*(1), 4-9.
- Ebesutani, C., Regan, J., Smith, A., Reise, S., Higa-McMillan, C., & Chorpita, B. F. (2012). The 10-item positive and negative affect schedule for children, child and parent shortened versions: application of item response theory for more efficient assessment. *Journal of Psychopathology and behavioral Assessment, 34*(2), 191-203.
- Eijssvogels, T. M., Thompson, P. D., & Franklin, B. A. (2018). The "Extreme Exercise Hypothesis": Recent Findings and Cardiovascular Health Implications. *Curr Treat Options Neurol 20*(10), 84.

- Eime, R. M., Young, J. A., Harvey, J. T., Charity, M. J., & Payne, W. R. (2013). A systematic review of the psychological and social benefits of participation in sport for children and adolescents: informing development of a conceptual model of health through sport. *IJBNPA*, 10(1), 98.
- Evenson, K. R., Catellier, D. J., Gill, K., Ondrak, K. S., & McMurray, R. G. (2008). Calibration of two objective measures of physical activity for children. *Journal of sports sciences*, 26(14), 1557-1565.
- Flegal, K. M., & Cole, T. J. (2013). *Construction of LMS parameters for the Centers for Disease Control and Prevention 2000 growth charts*: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics.
- Franco, B. G., & Cartiaux, O. (2016). Delta method and bootstrap in linear mixed models to estimate a proportion when no event is observed: application to intralesional resection in bone tumor surgery. *Statistics in medicine*, 35(20), 3563-3582.
- Goldfield, G. S., Henderson, K., Buchholz, A., Obeid, N., Nguyen, H., & Flament, M. F. (2011). Physical activity and psychological adjustment in adolescents. *J Phys Act Health*, 8(2), 157-163.
- Gutin, B., & Owens, S. (2011). The influence of physical activity on cardiometabolic biomarkers in youths: a review. *Ped Ex Sci*, 23(2), 169-185.
- Hamer, M., & Stamatakis, E. (2018). Relative proportion of vigorous physical activity, total volume of moderate to vigorous activity, and body mass index in youth: the Millennium Cohort Study. *IJO*, 1.
- Hay, J., Maximova, K., Durksen, A., Carson, V., Rinaldi, R. L., Torrance, B., . . . Veugelers, P. (2012). Physical activity intensity and cardiometabolic risk in youth. *Arch Pediatr Adolesc Med*, 166(11), 1022-1029.
- Janssen, I., & LeBlanc, A. G. (2010). Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *IJBNPA*, 7(1), 40.
- John, D., Sasaki, J. E., & Freedson, P. S. (2010). *Comparison of activity counts from the Actigraph GT3X and GT1M*. Paper presented at the International Congress on Physical Activity and Public Health (ICPAPH).
- Johnson, C. C., Murray, D. M., Elder, J. P., Jobe, J. B., Dunn, A. L., Kubik, M., . . . Schachter, K. (2008). Depressive symptoms and physical activity in adolescent girls. *MSSE*, 40(5), 818-826.
- Kremers, S. P., de Bruijn, G.-J., Droomers, M., van Lenthe, F., & Brug, J. (2007). Moderators of environmental intervention effects on diet and activity in youth. *American journal of preventive medicine*, 32(2), 163-172.
- Lagerberg, D. (2005). Physical activity and mental health in school children: A complicated relationship. *Acta Paediatrica*, 94(12), 1699-1701.

- Laurent, J., Catanzaro, S. J., Joiner Jr, T. E., Rudolph, K. D., Potter, K. I., Lambert, S., . . . Gathright, T. (1999). A measure of positive and negative affect for children: scale development and preliminary validation. *Psychological assessment*, 11(3), 326.
- Lonsdale, C., Lester, A., Owen, K. B., White, R. L., Moyes, I., Peralta, L., . . . MacMillan, F. (2015). An internet-supported physical activity intervention delivered in secondary schools located in Low socio-economic status communities: study protocol for the Activity and Motivation in Physical Education (AMPED) cluster randomized controlled trial. *BMC Public Health*, 16(1), 17.
- Lonsdale, C., Lester, A., Owen, K. B., White, R. L., Moyes, I., Peralta, L., . . . MacMillan, F. (2016). An internet-supported physical activity intervention delivered in secondary schools located in Low socio-economic status communities: study protocol for the Activity and Motivation in Physical Education (AMPED) cluster randomized controlled trial. *BMC Public Health*, 16(1), 17.
- Lubans, D. R., Plotnikoff, R. C., & Lubans, N. J. (2012). A systematic review of the impact of physical activity programmes on social and emotional well-being in at-risk youth. *Child and Adolescent Mental Health*, 17(1), 2-13.
- Masse, R., Poulin, C., Dassa, C., Lambert, J., Belair, S., & Battaglini, M. (1998). Elaboration and validation of a tool to measure psychological well-being: WBMMS. *Can J Public Health*, 89(5), 352-357.
- Mitchell, J. A., Dowda, M., Pate, R. R., Kordas, K., Froberg, K., Sardinha, L. B., . . . Page, A. (2017). Physical activity and pediatric obesity: a quantile regression analysis. *Medicine and science in sports and exercise*, 49(3), 466.
- Morgan, P. J., Young, M. D., Smith, J. J., & Lubans, D. R. (2016). Targeted health behavior interventions promoting physical activity: a conceptual model. *Exerc Sport Sci Rev.*, 44(2), 71-80.
- North, T. C., McCullagh, P., & Tran, Z. V. (1990). Effect of exercise on depression. *Exerc Sport Sci Rev.*, 18(1), 379-416.
- Owen, K. B., Parker, P. D., Astell-Burt, T., & Lonsdale, C. (2018). Effects of physical activity and breaks on mathematics engagement in adolescents. *Journal of science and medicine in sport*, 21(1), 63-68.
- Parfitt, G., Pavey, T., & Rowlands, A. V. (2009). Children's physical activity and psychological health: the relevance of intensity. *Acta Paediatrica*, 98(6), 1037-1043.
- Poitras, V. J., Gray, C. E., Borghese, M. M., Carson, V., Chaput, J.-P., Janssen, I., . . . Kho, M. E. (2016). Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. *Applied Physiology, Nutrition, and Metabolism*, 41(6), S197-S239.
- Raichlen, D. A., Foster, A. D., Seillier, A., Giuffrida, A., & Gerdeman, G. L. (2013). Exercise-induced endocannabinoid signaling is modulated by intensity. *Eur J Appl Physiol*, 113(4), 869-875.

- Rhodes, R. E., Janssen, I., Bredin, S. S., Warburton, D. E., & Bauman, A. (2017). Physical activity: Health impact, prevalence, correlates and interventions. *Psychology & Health*, 32(8), 942-975.
- Rhodes, R. E., & Kates, A. (2015). Can the affective response to exercise predict future motives and physical activity behavior? A systematic review of published evidence. *Annals of Behavioral Medicine*, 49(5), 715-731.
- Ryan, R. M., & Deci, E. L. (2001). On happiness and human potentials: A review of research on hedonic and eudaimonic well-being. *Annual review of psychology*, 52(1), 141-166.
- Spruit, A., Assink, M., van Vugt, E., van der Put, C., & Stams, G. J. (2016). The effects of physical activity interventions on psychosocial outcomes in adolescents: A meta-analytic review. *Clinical psychology review*, 45, 56-71.
- World Health Organization. (2004). *Promoting mental health: Concepts, emerging evidence, practice: Summary report*. Retrieved from
- World Health Organization. (2012). *Recommended levels of physical activity for children aged 5-17 years*. Retrieved from http://www.who.int/dietphysicalactivity/factsheet_young_people/en/
- Wu, X. Y., Han, L. H., Zhang, J. H., Luo, S., Hu, J. W., & Sun, K. (2017). The influence of physical activity, sedentary behavior on health-related quality of life among the general population of children and adolescents: A systematic review. *PloS one*, 12(11), e0187668.
- Zhang, Y., Zhang, D., Jiang, Y., Sun, W., Wang, Y., Chen, W., . . . Zhang, J. (2015). Association between physical activity and teacher-reported academic performance among fifth-graders in Shanghai: a quantile regression. *PloS one*, 10(3), e0115483.

Figure captions

Figure 1: Relationship between vigorous physical activity and affect in adolescents

Figure 2: Sex sensitivity analysis

Figure 3: Quartile regression sensitivity analysis

Table 1: Baseline characteristics of sample (n=1223)	
Age in years (mean(SD))	12.94 (0.54)
Sex (%)	
Male	55.1%
Female	44.9%
Body Mass Index (BMI) (mean(sd))	22.68 (4.99)
Physical Activity Intensity (mean(sd))	
Total light PA (mins/day)	81.51 (27.21)
Total moderate PA (mins/day)	27.95 (11.07)
Total vigorous PA (mins/day)	18.56 (10.91)
Daily accelerometer wear time (min/day) (mean(sd))	709.94 (120.38)
Affect (mean(sd))	
Positive affect (0-25)	18.05(4.52)
Negative affect (0-25)	9.33 (4.09)

Table 3: Estimates (β and Std. Error) of the association between positive and negative affect with physical activity intensity in the study sample

Predictors (Fixed-effects)	Positive affect			Negative affect		
	Un- standardized coefficients	p value	Standardized coefficients	Un- standardized coefficients	p value	Standardized coefficients
LPA (min/day)	0.047	0.274	0.300	0.001	0.837	0.010
LPA ² (min/day)	-0.000	0.262	-0.269	-0.000	0.197	-0.053
Mod (min/day)	-0.008	0.651	-0.0232	0.027	0.127	0.078
Mod ² (min/day)	-0.000	0.052	-0.096	-0.000	0.399	-0.037
Vig (min/day)	0.115	<0.001	0.307	-0.088	<0.001	-0.300
Vig ² (min/day)	-0.001	<0.001	-0.209	0.001	<0.001	0.184

All models were further controlled for age, BMI-z score at baseline, wear time, and each other physical activity intensity.

Table 4. Curvilinear effect of vigorous physical activity on well-being at different percentiles of the outcome (well-being) distribution

	Beta coefficient (linear/quadratic)			
	25 th Percentile	50 th Percentile	75 th Percentile	90 th Percentile
Positive affect	0.128/-0.001*	0.120/-0.001*	0.080/-0.000*	0.104/-0.000*
Negative affect	-0.080/0.001*	-0.108/0.001*	-0.102/0.001*	-0.081/0.001
* $p < 0.05$				

Relationship between VPA and Affect in Adolescents




Figure 1

Gender sensitivity analysis




Figure 2




Figure 3