NOVA

University of Newcastle Research Online

nova.newcastle.edu.au

Beaudry-Bellefeuille, Isabelle; Lane, Shelly J.; Ramos-Polo, Eduardo. "The Toileting Habit Profile Questionnaire: screening for sensory-based toileting difficulties in young children with constipation and retentive fecal incontinence." Published in the *Journal of Occupational Therapy, Schools, & Early Intervention* Vol. 9, Issue 2, p. 163-175 (2016).

Available from: <http://dx.doi.org/10.1080/19411243.2016.1141081>

This is an Accepted Manuscript of an article published by Taylor & Francis in the *Journal of Occupational Therapy, Schools, & Early Intervention* on 7 June 2016, available online: <https://www.tandfonline.com/doi/full/10.1080/19411243.2016.1141081>.

Accessed from: <http://hdl.handle.net/1959.13/1324027>

The Toileting Habit Profile Questionnaire: screening for sensory based toileting difficulties in young children with constipation and retentive fecal incontinence

Journal:	<i>Journal of Occupational Therapy, Schools, & Early Intervention</i>
Manuscript ID	WJOT-2015-0019.R1
Manuscript Type:	Original Article
Keywords:	Sensation disorders, Occupational Therapy, Constipation, Fecal incontinence, Toilet training

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Title: The Toileting Habit Profile Questionnaire: screening for sensory based toileting difficulties in young children with constipation and retentive fecal incontinence

Short title: The Toileting Habit Profile Questionnaire

Abstract

The Toileting Habit Profile Questionnaire: screening for sensory based toileting difficulties in young children with constipation and retentive fecal incontinence

Aims: Bowel management is an activity of daily living about which occupational therapists are often concerned. When problematic, bowel management issues can significantly limit participation in key occupations. Recently research has examined the relationship between sensory issues and bowel management concerns in children, indicating elimination problems occur more frequently in children with sensory processing difficulties than children without such problems. No tools currently provide information on toileting habits from a sensory perspective. This study examined the face and preliminary content validity of the Toileting Habit Profile Questionnaire, a tool designed to screen for sensory-based defecation difficulties in children.

Methods: A panel of experts (occupational therapists and gastroenterologists) reviewed a pilot version of the Toileting Habit Profile Questionnaire and responded to specific probe questions. Two reviewers conducted direct content analysis of responses; 100% agreement was reached on coding categories for narrative responses.

Results: Overall, experts demonstrated a high degree of agreement on the item content and intended use of the instrument. Input from the panel was useful in improving individual items and establishing this tool for initial research and clinical use.

Conclusions: The Toileting Habit Profile Questionnaire appears to capture behaviours that are consistent with problematic toileting behaviours, particularly as they relate to sensory over-reactivity.

Significance: The Toileting Habit Profile Questionnaire provides researchers and clinicians with a tool designed to identify the sensory issues that appear to be at the root of the toileting participation difficulties that many children with constipation and faecal incontinence experience.

Short abstract (100 words): This study examined the face and preliminary content validity of the Toileting Habit Profile Questionnaire, a tool designed to screen for sensory-based defecation difficulties in children. A panel of experts reviewed a pilot version of the questionnaire and responded to probe questions. Two reviewers conducted direct content analysis of responses; 100% agreement was reached. Experts demonstrated a high degree of agreement and their input was useful in establishing this tool for initial research and clinical use. The Toileting Habit Profile Questionnaire appears to capture behaviors that are consistent with problematic toileting behaviors, particularly as they relate to sensory over-reactivity.

Key words: constipation, fecal incontinence, occupational therapy, sensation disorders, toilet training

Occupational therapists are considered experts in understanding and supporting the everyday activities in which people engage, and this includes activities of daily living (World Federation of Occupational Therapists [WFOT], 2012). Bowel management comprises one important activity of daily living (American Occupational Therapy Association, 2014), and issues in this area can limit an individual's independence and social participation. Additionally, acquiring continence of bowel is considered an important milestone of childhood. As such, addressing issues related to bowel management is an important component of occupational therapy practice.

Fecal incontinence and constipation

Fecal incontinence is a common, under-treated and frequently misinterpreted disorder in children that often leads to impaired social acceptance, relationships and development (Friman, Hofstader, & Jones, 2006). It is one of the main symptoms of constipation, and constipation is one of the most common gastrointestinal complaints in children (Tabbers, Boluyt, Berger, & Benninga, 2011a). Worldwide prevalence of constipation is estimated to be 12% (range of 0.7 % to 29.6 %) in the general childhood population, with peak incidence occurring during toilet training (Mugie, Benninga, & Di Lorenzo, 2011). The term retentive fecal incontinence (RFI) is used to specify the presence of constipation and differentiate it from non-retentive fecal incontinence, a condition that occurs in the absence of stool retention. Children with RFI often soil due to rectal overflow, an involuntary process where soft stool from the bowel slips around a hard mass of stool that remains accumulated in the rectum (Cohn, 2011). This is highly problematic from an occupational perspective because the unpleasant odors caused by feces are often the cause of rejection and ridicule by peers, which can affect social participation and participation in key activities of childhood (Handley-More, Richards, Macauley, & Tierra, 2009).

1
2
3
4
5 Importantly, children with long lasting symptoms of constipation and fecal incontinence are at
6
7 risk for experiencing lower health-related quality of life (Kovacic et al., 2015). Children and
8
9 adolescents with fecal incontinence are at risk of experiencing peer related abuse and long lasting
10
11 psychosocial and emotional health problems (Kovacic et al., 2015). Parents also perceive lower
12
13 quality of life for themselves (Kovacic et al., 2015).
14
15

16
17 It is generally accepted that RFI is caused by either physical (e.g. a change in diet) or
18
19 psychological problems (e.g. birth of a sibling) (Cox et al., 2003). The event results in fecal
20
21 impaction of hard, large stools which may be painful and difficult to expulse. Subsequently
22
23 defecation may be anticipated as painful and/or difficult; the child may ignore the urge to
24
25 defecate and refuse to go to the toilet (Cox et al., 2003). Although the precipitating event leading
26
27 to RFI is not always clearly identified, the behavior of children with RFI seems to be partly
28
29 responsible for the development and/or maintenance of the condition. Stool withholding, possibly
30
31 due to pain (Cohn, 2011; Tabbers et al., 2011a), avoidance of using a toilet outside the home
32
33 (Tam et al., 2012), or stool toileting refusal/fear of sitting on the potty (Taubman, 1997) are
34
35 established as behaviors contributing to RFI. Clinically, occupational therapists have observed
36
37 that some children develop RFI in the absence of clear physical or psychological factors; they
38
39 appear to react in an exaggerated way to normal bodily sensations or certain aspects of toileting,
40
41 and come to anticipate defecation with a fearful response (Beaudry Bellefeuille & Ramos Polo,
42
43 2011; Beaudry, Schaaf, & Ramos, 2013).
44
45
46
47
48

49
50 **RFI and sensory processing**
51

52 It has been hypothesized that difficulty processing and integrating sensory information
53
54 could be a factor contributing to the development of some problematic behaviors in children
55
56 (Dunn, 2007; Hazen et al., 2008; Schaaf et al., 2010). Exaggerated responses to normal sensory
57
58
59
60

1
2
3
4 stimuli, sensory over-responsivity, have been associated with refusal to comply with parental
5
6 demands or atypical habits in relation to other types of self-care activities such as dressing or
7
8 feeding (Dunn, 2007; Hazen et al., 2008; Nadon, Ehrmann-Feldman, Dunn, & Gisel, 2011;
9
10 Schaaf et al., 2010). Occupational therapists working with children with RFI have suggested that
11
12 some of the above mentioned behaviors typical of children with RFI could be related to sensory
13
14 over-responsivity ([name deleted], 2011; [name deleted], 2013; Handley-More et al., 2009).
15
16 Recent research supports a relationship between sensory issues, fecal incontinence, constipation
17
18 and other gastrointestinal problems in children (Bakker, Boer, Benninga, Koelman, & Tijssen,
19
20 2010; Beaudry Bellefeuille & Ramos Polo, 2011; Beaudry et al., 2013; Beaudry-Bellefeuille,
21
22 2014; Mazurek et al., 2013; Pollock, Metz, & Barabash, 2014). Clearly defining these
23
24 relationships would lay the groundwork for an occupational performance approach to treatment
25
26 in conjunction with the more common medical management for children with RFI.
27
28
29
30
31

32
33 Assessment of the sensory responsivity basis for RFI is in its infancy. However,
34
35 conducting valid assessments to gain information about sensory processing factors that may be
36
37 affecting the child's participation in daily life is critical (Schaaf & Mailloux, 2015). Tools such as
38
39 the Sensory Profile (Dunn, 2014) or the Sensory Processing Measure (Parham, Ecker, Miller-
40
41 Kuhananek, Henry, & Glennon, 2007) examine the relationship between sensory responsivity
42
43 and the behaviors of young children during activities of daily living such as feeding and
44
45 grooming. However, currently no measure of sensory responsivity relative to participation in
46
47 bowel management habits is available for either clinical or research use.
48
49
50
51

52 The Toileting Habit Profile Questionnaire (THPQ) is a tool designed to address this gap,
53
54 screening for sensory based defecation difficulties. The THPQ was developed by an occupational
55
56 therapist in collaboration with a gastroenterologist. This team had observed that response to
57
58
59
60

sensations related to the evacuation of stool seemed to impact the acceptance of toilet training and the response to the urge to defecate in some children ([name deleted], 2011; [name deleted], 2013). The purpose of this study was to refine questions on the THPQ and examine its face and preliminary content validity.

The research questions posed were: 1) Do experienced pediatric gastroenterologists and occupational therapists working with young children with RFI consider the behaviors outlined in the THPQ to be characteristic of this population? ; 2) Do experienced pediatric gastroenterologists and occupational therapists working with young children with RFI consider the behaviors outlined in the THPQ to be related to issues in sensory reactivity?

Methods

Based on available literature and parent description of behaviors that are common to many children who are referred to OT for constipation and fecal incontinence, a clinical screening tool, the Toileting Habit Profile Questionnaire (THPQ), was developed. It has been used clinically for nearly 10 years and found to be highly useful in defining sensory concerns relative to constipation and fecal incontinence. In the current investigation the face and preliminary content validity of the THPQ was assessed through expert panel consultation.

Measure

The THPQ was developed in a bilingual (Spanish-English) format using simple and understandable wording (Grade level = 5.8 as determined through Microsoft WORD). The questionnaire is meant to be a screening tool to help differentiate typical defecation behaviors and reactions from those that are associated with constipation and fecal incontinence potentially related to sensory processing concerns. The original THPQ consisted of 11 questions, divided in

two sections: a) over-reactivity (9 questions) and b) under-reactivity (2 questions). Scored using a 5 point Likert scale (almost always through never), questions on the pre-study version of the TPHQ included the following:

1. My child hides while defecating.
2. My child asks for a diaper when he feels the need to defecate.
3. My child refuses to sit on the potty or the toilet to defecate.
4. My child always follows the same ritual when defecating.
5. My child seems to feel pain when defecating.
6. My child defecates only when paying attention to something else (while playing or watching television for example).
7. My child refuses to go to the toilet outside of the home.
8. My child's reaction to the odor of his/her feces is exaggerated.
9. My child refuses to wipe or be wiped after defecating.
10. My child does not seem to feel the urge to defecate.
11. My child does not realize he has soiled (feces) his clothes

Study design

This investigation assessed the face and preliminary content validity of the THPQ by requiring a panel of experts to respond to five probes relative to each THPQ item. The panel was comprised of six experts; three pediatric gastroenterologists with expertise in RFI, and three occupational therapists with expertise in Ayres Sensory Integration® and knowledge of potential issues related to activities of daily living, including toileting, in children. Consistent with the bilingual nature of the THPQ, all experts were bilingual (English-Spanish). The gastroenterologists represented private practice and the major hospitals of the [city name deleted].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The occupational therapists represented private practice and academia in diverse areas of [region name deleted]. No one on the panel were from the facility where the THPQ was developed and used clinically. Approval by the [name deleted] University Institutional Review Board was obtained before the beginning of the study.

Face validity, an initial step in test development, substantiates that the content of a tool appears to be measuring constructs of interest. Although a relatively weak reflection of validity, face validity is strengthened by approaching it systematically (Trochim & Donnelly, 2007). In the current study it reflects the judgement of the carefully selected experts from two professional fields. Content validity moves the process one step further, examining the relevance of each test item, and considering the relevance of each item to the overall construct being measured (Sireci & Sukin, 2013). Using panel members with both subject matter and experiential expertise strengthened this initial examination of content validity (Sireci & Sukin, 2013).

The THPQ and probe questions were distributed to the panel using the REDCap electronic data capture tools hosted at [name deleted] University (Harris, et al., 2009). As suggested by Sireci and Sukin (2013), probes asked experts to examine many elements of the THPQ, including item interpretation (How do you interpret what the item is asking?), potential reasons for behavior (Why do you think a child would have such a behavior?), likelihood of the behavior to be seen in typical children (Do you think that typically developing children have this behavior?), likelihood of the behavior in children with constipation and fecal incontinence (Do you think that this behavior is common in children with constipation and fecal incontinence?), and potential links to sensory over-responsivity (Do you think that this behavior could be related to over-responsivity to the sensations related to defecation (feel of potty/toilet on skin, anal/rectal distention, smell of feces, etc.))? Responses to the open ended questions were narrative. All

1
2
3
4
5 comments made by the respondents were taken into account. The feedback from the expert panel
6
7 was used to modify THPQ questions as needed. As suggested by Hsieh and Shannon (2005), both
8
9 an open-ended question (#2) and a targeted question (#5) were used to gain insight concerning
10
11 the relationship between the constructs of interest, in this case sensory responsivity and toileting
12
13 behaviors.
14

15
16 Direct content analysis methodology was used to establish coding of narrative responses
17
18 from the experts (Hsieh & Shannon, 2005). A direct content analysis approach is useful to
19
20 extend conceptually an existing theoretical framework and determine the initial coding scheme
21
22 (Hsieh & Shannon, 2005). Ayres Sensory Integration ® theory served as the theoretical
23
24 framework for this study. This framework considers emotional and behavioral reactions to the
25
26 sensory aspects of daily occupations a valid measure of sensory responsivity (Dunn, 2014;
27
28 Parham et al., 2007). As such, the responses of the experts on question #2 were coded in
29
30 accordance to manifestations hypothesized to be indicators of atypical sensory responsivity.
31
32 Based on Schaaf and Lane's (2014) review of sensory features, sensory over-responsivity is often
33
34 described in terms of negative emotional reactions to sensory stimuli such as fear or anxiety.
35
36 Further, avoiding, withdrawing, feeling pain or experiencing stress in relation to sensory stimuli
37
38 are common descriptors of sensory over-responsivity (Schaaf & Lane, 2014). Expressing distress,
39
40 dislike, repulsion or unusual and/or restricted preferences towards sensory stimuli, as well as
41
42 following rituals in personal hygiene are also considered signs of sensory over-responsivity
43
44 (Dunn, 2014). Diminished awareness, or lack of reaction to sensory stimuli, are common
45
46 descriptors of sensory under-responsivity (Schaaf & Lane, 2014). Responses not related to
47
48 sensory reactivity were classified together. Based on these theoretical guidelines, the authors
49
50 separately coded the experts' responses. Coding for question 1 (interpretation of question)
51
52
53
54
55
56
57
58
59
60

1
2
3
4 included “accurate”, “proposed reasons”, “inaccurate”, and “no response”. Coding for question 3
5
6 (is this behavior seen in typical children) included “yes”, along with qualifiers, and “no”. Coding
7
8 for questions 4 (is this behavior common in children with constipation and fecal incontinence)
9
10 and 5 (is this behavior related to over-reactivity to the sensations related to defecation) included
11
12 options for “yes”, “no”, and “don’t know”. Coding for question 2 included “sensory over-
13
14 responsivity”, “sensory under-responsivity”, and “other”; other included such things as cognitive
15
16 limitations and motor control concerns. Authors coded independently, reviewed coding together,
17
18 and engaged in discussion to reach agreement; 100% agreement was reached on all coding.
19
20
21
22

23
24 **Results**

25
26 Rather than offer their individual interpretation of the THPQ questions, requested by the
27
28 first probe question, “How do you interpret what the item is asking?”, experts generally proposed
29
30 reasons for the behaviors described by the items. One expert systematically left this question
31
32 blank. Responses to the remainder of the probe questions suggested that all questions were
33
34 accurately interpreted.
35
36

37
38 Responses to the second probe question, asking why such a behavior might be present,
39
40 were examined to gain insight concerning the relationship between sensory reactivity and
41
42 toileting behaviors. Experts frequently responded with more than one rationale, although general
43
44 agreement between experts was found. For all items, a majority of experts ($\geq 66.7\%$) considered
45
46 that sensory factors contributed to the behaviors described (Table 1).
47
48

49
50 *Insert Table 1 about here*
51

52
53 Probe question 3, “Do you think that typically developing children have this behavior?”,
54
55 revealed that many behaviors hypothesized to be associated with defecation difficulties may
56
57 occur, albeit infrequently, in typically developing children (Table 2).
58
59
60

Insert Table 2 about here.

The fourth probe question, “Do you think that this behavior is common in children with constipation and fecal incontinence?”, sought to examine the inclusion of items on the basis of frequency of occurrence. Responses showed high agreement among experts; over 90% of the items obtaining at least 66.7% agreement in the direction expected by the investigator (Figure 1). Item 8, “My child’s reaction to the odor of his/her feces is exaggerated”, although not considered a common behavior, was maintained on the revised version of the THPQ as there was 100% agreement on its relationship to sensory over-responsivity (probe question 5).

Insert Figure 1 about here.

Our final probe question, “Do you think that this behavior could be related to over-responsivity to the sensations related to defecation (feel of potty/toilet on skin, anal/rectal distention, smell of feces, etc.)?” showed high agreement in the expected direction on items 1,2,3,4,7,8,9 (Figure 2).

Insert Figure 2 about here.

The experts were divided when responding to the final probe question for items 5, 6, 10 and 11. On item 5 (“My child seems to feel pain when defecating”), although there was high agreement in manifestations of sensory over-responsivity in the narrative responses of probe question number 2, only half of the experts indicated that pain felt while defecating could be related to sensory over-responsivity. Amongst those who felt it was not related, two clarified that feeling pain was probably due to the hard consistency of feces.

Responses concerning item 6 (“My child defecates only when paying attention to something else.”) included three supporting a relationship with over-responsivity, two indicating this was not related, and one expert did not respond. A review of our recent clinical data on this

item has shown that it is neither common nor related to over-responsivity. Considering this information, item 6 was eliminated from the revised version of the THPQ.

Items 10 (“My child does not seem to feel the urge to defecate.”) and 11 (“My child does not realize he has soiled (feces) his clothes.”) were included on the THPQ to reflect under-responsivity to sensory input; experts showed 100% agreement in this direction in their narrative responses to question 2. When experts were asked “Do you think that this behavior could be related to over-responsivity to the sensations related to defecation?” it was expected that they would unanimously respond *no*; not feeling the urge to defecate or not noticing that one is soiled with feces is not a behavior that is expected to be related to sensory over-responsivity. Surprisingly some experts responded affirmatively to probe question 5 for items 10 and 11.

Conclusions

Face and preliminary content validity of the THPQ was established in this investigation using a panel of experts consisting of pediatric gastroenterologists and occupational therapists with expertise in Ayres Sensory Integration®. Based on directed content analysis we found a high degree of agreement among the members of the expert panel on the relevancy of items and their relationship to sensory responsivity. Findings were considered relative to existing clinical experience, and some changes have been incorporated into the THPQ, noted below, improving this tool before its use in a pilot study and in clinical practice.

Item 5 was not fully clear to all panel members. Our clinical experience shows that many children continue to manifest pain even when they take medications which soften stools. We hypothesize that feeling pain while defecating could be due to over- responsivity to typical bodily

1
2
3
4 sensations. Item 5 was modified to include a reference to the consistency of the stool; “My child
5
6 seems to feel pain when defecating even if the stool is soft.”
7
8

9
10 Experts agreed that Items 10 and 11 tapped into sensory under-responsivity; this was as
11
12 expected. However, such behaviors are not usually present in children who present with over-
13
14 responsiveness, yet some experts indicated they might be identified. These questions were
15
16 maintained in the revised version of the THPQ, and incompatible responses between the first
17
18 section of the THPQ (over-responsivity) and the last 2 items (under-responsivity) would alert the
19
20 examiner to possible misunderstanding by the respondent. For example, clinical experience
21
22 shows that some children with sensory over-responsivity ignore or deny that they are soiled for
23
24 fear of punishment or ridicule and this behavior may be interpreted as under-responsivity.
25
26 Detailed observation and interview with caregivers is needed to fully understand the child’s
27
28
29
30
31 behavior.
32

33 The fact that children may fluctuate in their responses to sensation or manifest over-
34
35 responsiveness to some sensations and under-responsivity to others must also be considered
36
37 (Baranek, David, Poe, Stone, & Watson, 2006; Kientz & Dunn, 1997; Parham & Mailloux,
38
39 1996). However in the case of children with retentive fecal incontinence who do not respond to
40
41 conventional medical management, difficulties accepting toileting and defecation appear to be
42
43 related mainly to over-responsivity in the tactile system; normal sensations such as the passage of
44
45 feces or the contact of the potty/toilet on the skin are interpreted as painful. Further research is
46
47
48
49
50 needed to clarify the sensory response patterns of children with different types of defecation
51
52 issues.
53
54
55
56
57
58
59
60

Based on content analysis of these expert responses, the revised version of the THPQ includes 10 items; the first 8 items are hypothesized to be related to sensory over-responsivity and items 9 and 10 to sensory under-responsivity.

1. My child hides while defecating.
2. My child asks for a diaper when he feels the need to defecate.
3. My child refuses to sit on the potty or the toilet to defecate.
4. My child always follows the same ritual when defecating.
5. My child seems to feel pain when defecating, even if the stool is soft.
6. My child refuses to go to the toilet outside of the home.
7. My child’s reaction to the odor of his/her feces is exaggerated.
8. My child refuses to wipe or be wiped after defecating.
9. My child does not seem to feel the urge to defecate.
10. My child does not realize he has soiled (feces) his clothes.

The construct of sensory under and over-responsivity is well established (Dunn, 2014; Lane, Reynolds, & Thacker, 2010; Parham et al., 2007; Reynolds, Bendixen, Lawrence, & Lane, 2011; Su & Parham, 2014). Caregiver questionnaires are available to systematically document behavioral manifestations of passiveness, discomfort, distress or intolerance in relation to daily occupations which include specific sensory input (Dunn, 2014; Parham et al., 2007). For example expressing distress in relation to grooming is considered a sign of tactile over-responsivity and may be a factor related to a child’s difficulty participating in dressing and bathing. However, these tools do not include items related to toileting, a crucial childhood occupation. Such items could be useful to better understand difficulties participating in toileting routines, a frequent complaint of parents of children with retentive fecal incontinence.

1
2
3
4
5 The THPQ has been used clinically to screen for sensory processing concerns in children
6
7 with RFI that has not responded to first line medical management. Developed based on careful
8
9 documentation of the behaviors and the progress of the children referred for possible sensory
10
11 processing deficits relative to RFI, the THPQ appears to capture behaviors that are consistent
12
13 with problematic toileting behaviors, particularly as they relate to sensory over-reactivity ([name
14
15 deleted], 2011; [name deleted] , 2014).
16
17

18
19 This study was part of a larger study in which toileting behaviors reported by parents of
20
21 typically developing children were shown to be significantly different from those reported by
22
23 parents of children with RFI ([name deleted], 2014). This finding supports the discriminative
24
25 validity of the THPQ. Interestingly, sensory responsivity as measured by the Spanish version of
26
27 the Short Sensory Profile was also significantly different between the two groups, suggesting that
28
29 the behaviors described in the THPQ may be useful to identify sensory based defecation issues
30
31 ([name deleted], 2014). Further investigation is warranted.
32
33
34

35
36 There is growing evidence that OT is successful in improving participation in children
37
38 with sensory issues (Pfeiffer, Koenig, Kinnealy, Sheppard, & Henderson , 2011; Schaaf, 2011;
39
40 Schaaf et al., 2013). There is also some evidence that addressing the sensory issues that appear to
41
42 be at the root of the behaviors related to the development and maintenance of constipation and
43
44 fecal incontinence may contribute to more successful treatment outcomes for children who
45
46 experience this complex and often chronic condition ([name deleted], 2011; [name deleted],
47
48 2013; Handley-More et al., 2009). The THPQ may provide a structure useful in identifying and
49
50 tracking these behaviors over the course of treatment.
51
52
53
54
55
56
57
58
59
60

Limitations and Future Research

The main limitation to this study was the relatively small size of the expert panel. However, having bilingual experts with substantial clinical experience in pediatric gastroenterology and Ayres Sensory Integration was seen as a strength in establishing face and content validity. Additionally, although feedback from the parents was obtained during the development of the THPQ and during a preliminary study using the tool, formal consultation on face validity did not include parents. Future research will need to strengthen the validity of this tool and include the consumer perspective.

Acknowledgements

Our gratitude to the experts who kindly took the time to review the THPQ. Many thanks to [name deleted], experienced translation consultant, for his review of the pre-study version of the THPQ.

References

- American Occupational Therapy Association. (2014). Occupational therapy practice framework: Domain and process (3 rd ed.). *American Journal of Occupational Therapy*, 68(Suppl. 1), S1-S48. doi:10.5014/ajot.2014.682006.
- Bakker, M. J., Boer, F., Benninga, M. A., Koelman, J. H., Tijssen, M. A. (2010). Increased auditory startle reflex in children with functional abdominal pain. *Journal of Pediatrics*, 156(2), 285-291.
- Baranek, G. T., David, F. J., Poe, M. D., Stone, W. L., & Watson, L. R. (2006). Sensory experience questionnaire: Discriminating sensory features in young children with autism, developmental delays, and typical development. *Journal of Child Psychology and Psychiatry*, 47, 591–601.
- Beaudry Bellefeuille, I., & Ramos Polo, E. (2011). Tratamiento combinado de la retención voluntaria de heces mediante fármacos y terapia ocupacional [Combined treatment of voluntary stool retention with medication and occupational therapy]. *Boletín de la Sociedad de Pediatría de Asturias, Cantabria, Castilla y León*, 51, 169-176.
- Beaudry, I. B., Schaaf, R. C., & Ramos, E. P. (2013). Brief Report—Occupational therapy based on Ayres Sensory Integration in the treatment of retentive fecal incontinence in a 3-year-old boy. *American Journal of Occupational Therapy*, 67, 601–606.
<http://dx.doi.org/10.5014/ajot.2013.008086>
- Beaudry-Bellefeuille, I. (2014). Examining the Sensory Characteristics of Preschool Children with Retentive Fecal Incontinence. VCU Theses and Dissertations. Paper 3327.
<http://scholarscompass.vcu.edu/etd/3327>
- Cohn, A. (2011). Clinical features, psychological issues and management of constipation in childhood. *Nursing Children and Young People*, 23(3), 29-35.
- Cox, D. J., Ritterband, L. M., Quillian, W., Kovatchev, B., Morris, J., Sutphen, J., Borowitz, S. (2003). Assessment of behavioral mechanisms maintaining encopresis: Virginia

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

encopresis-constipation apperception test. *Journal of Pediatric Psychology* 28(6): 375-382.

Dunn, W. (2014). *Sensory Profile™ 2*. San Antonio, TX: Pearson Education.

Dunn, W. (2007). Supporting children to participate successfully in everyday life by using sensory processing knowledge. *Infants & Young Children*, 20(2), 84-101.

Friman, P. C., Hofstader, K. L., & Jones, K. M. (2006). A biobehavioral approach to the treatment of functional encopresis in children. *Journal of Early and Intensive Behavior Intervention*, 3(3), 263-271.

Handley-More, D., Richards, K., Macauley, R., & Tierra, A. (2009). Encopresis: Multidisciplinary management. *Journal of Occupational Therapy, Schools, & Early Intervention*, 2(2), 96–102. doi: 10.1080/19411240903146400

Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research Electronic Data Capture (REDCap) - A metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics*, 42(2), 377–381. doi:10.1016/j.jbi.2008.08.010

Hazen, E. P., Reichert, E. L., Piacentini, J. C., Miguel, E., Do Rosario, M., Pauls, D., & Geller, D. A. (2008). Case series: Sensory intolerance as a primary symptom of pediatric OCD. *Annals of Clinical Psychiatry*, 20(4), 199-203. doi:10.1080/10401230802437365

Hseih, H. F., Shannon, S. E. (2005). Three approaches to qualitative content analysis. *Qualitative Health Research* 15(9): 1277-1288. doi: 0.1177/1049732305276687

Kientz, M. A., & Dunn, W. (1997). A comparison of the performance of children with and without autism on the sensory profile. *American Journal of Occupational Therapy*, 51, 530-537.

Kovacic, K., Sood, M. R., Mugie S., Di Lorenzo, C., Nurko, S., Heinz, N., ..., Silverman, A. H. (2015). A multicenter study on childhood constipation and fecal incontinence: effects on

quality of life. *Journal of Pediatrics*, 166(6):1482-1487.e1. doi:
10.1016/j.jpeds.2015.03.016.

- Lane, S. J, Reynolds, S., & Thacker, L. (2010). Sensory over-responsivity and ADHD: Differentiating using electrodermal responses, cortisol, and anxiety. *Frontiers in Integrative Neuroscience*, 29(4), article 8. doi: 10.3389/fnint.2010.00008.
- Mazurek, M. O., Vasa, R. A., Kalb, L. G., Kanne, S., Rosenberg, D., Keefer, A., ... Lowery, L. A. (2012). Anxiety, sensory over-responsivity, and gastrointestinal problems in children with autism spectrum disorders. *Journal of Abnormal Child Psychology*. Advance online publication. doi 10.1007/s10802-012-9668-x
- Mugie, S. M., Benninga, M. A., Di Lorenzo C. (2011). Epidemiology of constipation in children and adults: A systematic review. *Best Practice & Research Clinical Gastroenterology*, 25(1), 3-18.
- Nadon, G., Ehrmann-Feldman, D., Dunn, W., & Gisell, E. (2011). Association of sensory processing and eating problems in children with autism spectrum disorders. *Autism Research and Treatment*. doi:10.1155/2011/541926
- Parham, D. L., Ecker, C., Miller-Kuahanneck, H., Henry, D. A., & Glennon, T. (2007). *Sensory Processing Measure (SPM) manual*. Los Angeles: Western Psychological Services.
- Parham, D. L., & Mailloux, Z. (1996). Sensory integration. In J. Case-Smith, A. S. Allen, & P. N. Pratt (Eds.), *Occupational Therapy for Children* (3rd ed., pp. 307–355). St Louis: Mosby.
- Pfeiffer, B. A., Koenig, K., Kinnealey, M., Sheppard, M., & Henderson, L. (2011). Effectiveness of sensory integration interventions in children with autism spectrum disorders: A pilot study. *The American Journal of Occupational Therapy*, 65(1), 76–85.
- Pollock, M. R., Metz, A. E., & Barabash, T. (2014). Brief Report—Association between dysfunctional elimination syndrome and sensory processing disorder. *American Journal of Occupational Therapy*, 68, 472–477. [http://dx.doi.org/ 10.5014/ajot.2014.011411](http://dx.doi.org/10.5014/ajot.2014.011411)

Reynolds, S., Bendixen, R. M., Lawrence, T., & Lane, S. J. (2011). A pilot study examining activity participation, sensory responsiveness, and competence in children with high functioning autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 41(11), 1496-1506. doi 10.1007/s10803-010-1173-x

Schaaf, R. C. (2011). Interventions that address sensory dysfunction for individuals with autism spectrum disorders: Preliminary evidence for the superiority of sensory integration compared to other sensory approaches. In B. Reichow, P. Doehring, D. V. Cicchetti, & F. R. Volkmar (Eds.), *Evidence-based practices and treatments for children with autism* (pp. 245–273). New York: Springer.

Schaaf, R. C., Benevides, T., Blanche, E., Brett-Green, B., Burke, J., Cohn, E., ... Schoen, S. (2010). Parasympathetic functions in children with sensory processing disorder. *Frontiers in Integrative Neuroscience*, 4(4). doi:10.3389/fnint.2010.00004

Schaaf, R. C., Benevides, T., Mailloux, Z., Faller, P., Hunt, J., van Hooydonk, E., ... Kelly, D. (2013). An intervention for sensory difficulties in children with autism: A randomized trial. *Journal of Autism and Developmental Disorders*. doi: 10.1007/s10803-013-1983-8

Schaaf, R. C., Lane, A. E. (2014). Toward a best-practice protocol for assessment of sensory features in ASD. *Journal of Autism and Developmental Disorders*. doi: 10.1007/s10803-014-2299-z

Schaaf, R. C., Mailloux, Z. (2015). *Clinician's Guide for Implementing Ayres Sensory Integration: Promoting Participation for Children with Autism*. Bethesda, MD: AOTA Press

Sireci, S.G., & Sukin, T. Test validity. In K.F. Geisinger, B.A. Bracken, J.F. Carlson, J-I.C. Hansen, N.R. Kuncel, S.P. Reise, & M.C. Rodriguez. (Eds), *APA handbook of testing and assessment in psychology*, Vol. 1: Test theory and testing and assessment in industrial and organizational psychology (pp. 61-84). Washington, DC, US: American Psychological Association.

- 1
2
3
4 Su, C. T., Parham, D. (2014). Validity of Sensory Systems as Distinct Constructs. *American*
5
6 *Journal of Occupational Therapy*, 68(5), 546-554. doi: 10.5014/ajot.2014.012518.
7
8
9 Tabbers, M. M., Boluyt, N., Berger, M. Y., & Benninga, M. A. (2011a). Diagnosis and treatment
10 of functional constipation. *European Journal of Pediatrics*, 170, 955–96.
11
12 doi:10.1007/s00431-011-1515-5.
13
14
15 Tam, Y. H., Li, A. M., So, H. K., Shit, K. Y., Pang, K.K., Wong, Y.S., ..., Lee, K. H. (2012).
16 Socioenvironmental factors associated with constipation in Hong Kong children and
17 Rome III criteria. *Journal of Pediatric Gastroenterology and Nutrition*, 55(1), 56-61.
18
19
20
21 Taubman, B. (1997). Toilet training and toileting refusal for stool only: A prospective study.
22 *Pediatrics*, 99(1), 54-58.
23
24
25
26 Trochim, W. M. K., & Donnelly, J. P. (2007). Research methods knowledge base, 3e.
27 Independence, KY: Cengage Learning Atomic Dog Publishing.
28
29
30 World Federation of Occupational Therapists [WFOT], (2012). *Position Statement. Activities of*
31 *Daily Living*. Forrestfield, Western Australia: WFOT.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Expert panel responses to probe question 2: Why do you think a child would have such a behavior?

Item	Why do you think a child would have such a behavior?
1. My child hides while defecating.	<ul style="list-style-type: none">• SOR : 5 (83.3 % *)• OTHER: 4 (Embarrassment: 3; Behavioral issue:1)
2. My child asks for a diaper when he feels the need to defecate.	<ul style="list-style-type: none">• SOR: 4 (66.7 % *)• OTHER:2 (Over-concious:1, Poor toilet training: 1)
3. My child refuses to sit on the potty or the toilet to defecate.	<ul style="list-style-type: none">• SOR:6 (100 % *)• OTHER:2 (Motor/cognitive issues: 1, over-conscious: 1)
4. My child always follows the same ritual when defecating.	<ul style="list-style-type: none">• SOR:5 (83.3 % *)• OTHER: 1 (behavioral/emotional disorders)
5. My child seems to feel pain when defecating.	<ul style="list-style-type: none">• SOR: 5 (83.3 % *)• OTHER: 1 (constipation)
6. My child defecates only when paying attention to something else.	<ul style="list-style-type: none">• SOR: 4 (66.7 % *)• SUR: 1• OTHER: 1 (TV as a reward)
7. My child refuses to go to the toilet outside of the home.	<ul style="list-style-type: none">• SOR: 5 (83.3 % *)• OTHER: 1 (associated with parental behavior)
8. My child’s reaction to the odor of his/her feces is exaggerated.	<ul style="list-style-type: none">• SOR: 5 (83.3 % *)• OTHER: 1 (Behaviorally reinforced response)
9. My child refuses to wipe or be wiped after defecating.	<ul style="list-style-type: none">• SOR: 5 (83.3 % *)• OTHER: 2 (motor impairments: 1, behavior:2)
10. My child does not seem to feel the urge to defecate.	<ul style="list-style-type: none">• SUR: 6 (100 % *)• OTHER: 1 (developmental delay.)
11. My child does not realize he has soiled (feces) his clothes.	<ul style="list-style-type: none">• SUR: 6 (100 % *)• OTHER: 2 (ignores: 2, intellectual disability:1)

Table note: * percent agreement among experts; numbers refer to number of responses (in some cases experts gave more than one rationale); SOR (sensory over-responsivity): fear, anxiety, pain, stress, repulsion, avoiding, withdrawing, dislike, unusual and/or restricted preferences towards sensory stimuli, following rituals in personal hygiene; SUR (sensory under-responsivity): diminished awareness, lack of reaction.

Table 2: Expert panel responses to probe question 3: Do you think that typically developing children have this behavior?

Item	Do you think that typically developing children have this behavior?
1 My child hides while defecating.	<ul style="list-style-type: none"> • Yes, but infrequent: 4# (66.7 % *) • No: 2#
2 My child asks for a diaper when he feels the need to defecate.	<ul style="list-style-type: none"> • Yes, but infrequent: 3# (50 % *) • No: 3#
3 My child refuses to sit on the potty or the toilet to defecate.	<ul style="list-style-type: none"> • Yes, but infrequent: 5# (83.3 % *) • No: 1#
4 My child always follows the same ritual when defecating.	<ul style="list-style-type: none"> • Yes: 1# • Yes, but infrequent: 2# • No: 3# (50 % *)
5 My child seems to feel pain when defecating.	<ul style="list-style-type: none"> • Yes, if constipated: 3# (50 % *) • Yes: 2# • No answer: 1#
6 My child defecates only when paying attention to something else.	<ul style="list-style-type: none"> • Yes but infrequent: 2# • No: 4 # (66.7 % *)
7 My child refuses to go to the toilet outside of the home.	<ul style="list-style-type: none"> • Yes: 4# (66.7 % *) • Yes but infrequent: 1# • No: 1#
8 My child's reaction to the odor of his/her feces is exaggerated.	<ul style="list-style-type: none"> • Yes: 2# • Yes but infrequent: 2# • No: 1# • No answer: 1#
9 My child refuses to wipe or be wiped after defecating.	<ul style="list-style-type: none"> • Yes but infrequent: 3# (50 % *) • No: 3#
10 My child does not seem to feel the urge to defecate.	<ul style="list-style-type: none"> • Yes: 1# • No: 4 # (66.7 % *) • No answer: 1#
11 My child does not realize he has soiled (feces) his clothes.	<ul style="list-style-type: none"> • Yes: 1# • Yes but infrequent: 2# • No: 3# (50 %*)

Table note: # number of respondents; * percent agreement

Figure 1

Figure 2