NOVA

University of Newcastle Research Online

nova.newcastle.edu.au

White, Rhiannon Lee; Babic, Mark J.; Parker, Philip D.; Lubans, David R.; Astell-Burt, Thomas; Lonsdale, Chris; "Domain-specific physical activity and mental health: a meta-analysis".
Published in American Journal of Preventive Medicine Vol. 52, Issue 5, p. 653-666 (2017)

Available from: <http://dx.doi.org/10.1016/j.amepre.2016.12.008>

© 2017. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Accessed from: <http://hdl.handle.net/1959.13/1354810>

Title

Domain-specific physical activity and mental health: A meta-analysis.

Authors and Affiliations

Rhiannon Lee White, Institute for Positive Psychology and Education, Australian Catholic University

Mark. J. Babic, Priority Research Centre for Physical Activity and Nutrition, University of Newcastle

Philip. D. Parker, Institute for Positive Psychology and Education, Australian Catholic University

David. R. Lubans, Priority Research Centre for Physical Activity and Nutrition, University of Newcastle

Thomas Astell-Burt, School of Health and Society, University of Wollongong

Chris Lonsdale*, Institute for Positive Psychology and Education, Australian Catholic University

***Corresponding author**

Associate Professor Chris Lonsdale, Locked Bag 2002, Strathfield, NSW, 2135, Australia, +612 9701 4642, Chris.Lonsdale@acu.edu.au

Abstract

Context: The mental health benefits of physical activity are well established. However, less is known about whether the relationship between physical activity and mental health is consistent across different life domains. It is important to understand how context may influence the relationship between physical activity and mental health so that interventions and policy guidelines can be tailored to maximise positive effects.

Evidence acquisition: In 2015, systematic searches of four databases identified 13,435 records, of which 98 studies met the inclusion criteria.

Evidence synthesis: Included studies were published between 1988 and 2015 and had a combined sample size of 648,726. Of the 98 studies included, 93 examined leisure-time physical activity, 14 examined work-related physical activity, 15 examined transport physical activity, 16 examined household physical activity, 3 examined school sport, and 3 examined physical education. Multi-level meta-analyses showed that leisure-time physical activity ($r = .13$) and transport physical activity ($r = .13$) both have a positive association with mental health. Leisure-time PA ($r = -.11$) and school sport ($r = -.09$) both have an inverse association with mental ill-health. However, physical activity is not consistently associated with lower mental ill-health across domains as work-related physical activity is positively associated with mental ill-health ($r = .09$). Household physical activity and participation in physical education had no relationship with mental health or mental ill-health.

Conclusions: The domain in which physical activity occurs influences the relationship between physical activity and mental health and should, therefore, be considered when developing interventions, treatment programs, and policy guidelines.

Context

Mental ill-health accounts for a substantial proportion of the burden of disease worldwide.¹⁻³ Individuals with mental ill-health may experience low self-esteem, struggle to maintain interpersonal relationships, and have a higher risk of communicable and non-communicable diseases than those not experiencing mental ill-health.^{1,4} Mental health, however, is a positive state of well-being where individuals realize their potential, experience positive emotions, and are able to cope with stress, maintain interpersonal relationships, work productively, and contribute to their community.⁵

Systematic reviews have shown that physical activity (PA) is associated with greater mental health,⁶ and a reduced risk of mental ill-health, specifically depression and anxiety.^{7,8} However, many of the studies included in these reviews measured total weekly PA. Given that PA is defined as any muscular movement that expends energy, PA can take place during leisure-time, work, or school, as a method of transportation, or while carrying out household chores.⁹ As the reasons for participation are likely to vary between these different PA domains (i.e., areas of life which present opportunities for PA), it is possible that the outcomes experienced are also different.^{10,11}

A literature review that compared total PA (i.e., all PA undertaken in a given time regardless of life domain) to leisure-time PA reported that leisure-time PA was more consistently and strongly associated with reduced depression.¹² Further, a number of individual studies have shown that leisure-time PA has a stronger relationship with mental health and mental ill-health, compared to work-related PA, transport PA, and household PA.¹³⁻¹⁵ In order to understand how to optimally use PA to promote mental health and prevent mental ill-health, an understanding of the relationships within specific PA domains is required. Therefore, the

primary purpose of this study was to synthesise study results and provide meta-analytic evidence of the relationships between domain-specific PA and mental health and mental ill-health. The secondary purpose was to conduct moderated meta-analyses to explore factors that may contribute to variation. The final purpose was to identify gaps in the literature regarding PA domains that have not been investigated extensively.

Evidence Acquisition

The methods detailed below were conducted in accordance with the ‘Preferred Reporting Items for Systematic Reviews and Meta-Analysis’ (PRISMA) statement.¹⁶

Inclusion and Exclusion Criteria

Studies eligible for this review met the following inclusion criteria: a) a quantitative assessment of PA within at least one specified domain (e.g., leisure-time, work, physical education, active commuting, or household or domestic duties), unless an experimental study was carried out within a specific domain then no domain specific measure of PA was required, b) a quantitative assessment of at least one mental health outcome variable (i.e., mental health, mental wellbeing, psychological wellbeing, subjective wellbeing, life satisfaction, positive affect, negative affect, depression, anxiety, stress, or psychological distress), c) a quantitative assessment of the relationship between PA in a specified domain and mental health, d) a cross sectional, longitudinal, or experimental design, and e) full text, peer reviewed journal articles.

With regards to the mental health related search terms, all studies measuring mental health, mental wellbeing, and psychological wellbeing were included.¹⁷ Further, positive affect and

life satisfaction are two core components of mental wellbeing and were also included.¹⁸ Mental ill-health is defined as a broad term encompassing mental health disorders and preclinical mental health problems.¹⁹⁻²² Internalising disorders are the largest cluster of mental disorders within the DSM-V and are responsible for a large portion of the burden of disease attributable to mental ill-health.^{2,23-25} Internalising disorders reflect the tendency to express mental distress inwards emotionally, and include unipolar depression and anxiety disorders.^{23,26} As such, mental ill-health in this review refers to symptoms of internalising mental disorders (i.e., depression and anxiety) and mental health problems (i.e., psychological distress, stress, negative affect).

Studies were excluded if they: a) included participants from special populations (e.g., participants who were physically frail), b) only measured total PA or did not specify the domain in which PA was measured, c) measured sedentary behaviour but not levels of PA, d) measured mental health specifically in terms of a particular setting or circumstance (e.g., job strain), e) solely reported qualitative data, or f) were published in languages other than English.

Search Strategy

Systematic searches were conducted in February 2015 in four databases: Scopus, PubMed, PsychINFO, and SPORTDiscus. Searches involved keywords from three main groups, including physical activity terms, mental health-related terms, and domain specific terms. The search was not restricted by publication date or study design, but included exclusion terms to reduce the number of irrelevant records identified. The full search strategy can be found in Table S1. Search results were then exported into Endnote reference manager software and removed duplicates.

Study Selection

Two reviewers independently assessed the title and abstract of each study and records recommended for exclusion by both reviewers were excluded. The same two reviewers then independently screened the full text of the remaining studies recommending each for inclusion or exclusion. Discrepancies were resolved by discussion between the two reviewers, along with a third reviewer, until consensus was reached. Finally, the reference lists of all included studies were checked for any relevant articles not returned in the search results.

Data Extraction

A researcher extracted data from each article into a predefined data extraction form. Data items included: authors, year of publication, study design, sample (size, age, sex, and country), measures of PA and mental health or mental ill-health, method of data analysis, and statistical results. A second reviewer cross-checked the extracted data for accuracy and all discrepancies were discussed and resolved.

Statistical Analysis

Correlation effect sizes were either extracted, or calculated from the results of each individual study.^{27,28} Next, internal consistency scores (e.g., Cronbach's α) were used to correct each correlation for attenuation.^{29,30} When this information was missing, an internal consistency score reported in similar literature was used. If an internal consistency score was not obtainable, or the study used a single item measure, a conservative score of .70 was used.³¹ Finally, because r becomes more skewed as the value moves away from zero, all r were

converted to Fisher's adjusted Z scale (z), which is almost normally distributed.^{27,28} The variance (v) and standard error (SE) of each effect size were then calculated.²⁷

Two random-effects meta-analyses (i.e., one for mental health and one for mental ill-health) were conducted in late 2015. In order to account for dependent effect sizes (i.e., multiple effect sizes within a single study), the metaSEM package in R, version 3.2.2, was used to conduct three-level meta-analyses using structural equation modelling, where multiple effect sizes (Level 2) are clustered within studies (Level 3).^{32 33} The weighted average effect sizes (r) and 95% CIs have been reported for each. In line with Cohen's guideline for interpreting the strength of correlations (r), a correlation of .1 was interpreted as weak, .3 moderate, and .5 strong.³⁴

The I^2 statistic indicates the percentage of the total variation in effect sizes that is due to genuine differences between the individual study results as opposed to chance.³⁵ However, as I^2 increases rapidly towards 100% as the number of participants within the primary studies increases, I^2 is interpreted as possible heterogeneity that may be relevant.³⁶ According to Cochrane's overlapping criteria, an I^2 statistic between 0% and 40 % may not be important, 30% to 60% may represent moderate heterogeneity, 50% to 90% may represent substantial heterogeneity, and >75% considerable heterogeneity.³⁷

Next, mixed-effects meta-analyses were conducted as they extend the random-effects meta-analyses by including fixed and random components within the one model in order to test study characteristics as potential predictors.³³ Study characteristics tested as potential moderators included PA domain, the specific mental health outcomes, age, sex, study design,

type of PA measure, and risk of bias. Age was categorised as children (1-9 years), adolescents (10-19 years), adults (20-64 years), and older adults (65+ years).^{38,39} For each mixed-effects meta-analysis the regression coefficient (β) is reported, demonstrating whether the moderator is a positive or negative predictor of the effect size. The R^2 which demonstrates the proportion of heterogeneity explained by the moderator variable is also reported.³² The p value for an omnibus ANOVA test demonstrating whether the mixed-effects model was significantly different from the random-effects model is also shown in Table 1 and Table 2. When the mixed-effects meta-analysis was significantly different, the effect size and 95% CI within each sub-group were included.

Rosenthal's fail-safe N was calculated to indicate the number of unpublished studies with a mean effect of zero required to cause the meta-analysis effect to become non significant ($p > .05$).⁴⁰ A larger fail-safe N , relative to the number of included studies, suggests that unpublished studies not included in the review would be unlikely to overturn the effect reported.⁴⁰ In order to assess the risk of bias across studies, Egger's regression asymmetry tests were conducted on the two overall meta-analyses (i.e., mental health and mental-ill health). However, it was not appropriate to conduct Egger's tests for the meta-analyses on each PA domain as ≥ 10 studies are required to distinguish chance from real asymmetry.^{37,41}

Risk of Bias of Individual Studies

Six criterion were adapted from the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement⁴² and the Consolidated Standards of Reporting Trials (CONSORT) statement to assess risk of bias of individual studies.⁴³ The criteria used was similar to that used in previous meta-analyses.^{44,45} Two reviewers independently assigned a 1 (present and explicitly explained) or a 0 (absent or inadequately described) to each included

study on each criterion: a) random selection of schools/and or participants (sampling procedures were appropriate and adequately described, including randomisation process for experimental studies), b) participant eligibility criteria stated and adequate description of study sample, c) valid assessment of PA participation including adherence to a PA program if no direct measure of PA was included, d) valid assessment of mental health or mental ill-health, e) covariates adjusted for in analyses relating to PA and mental health (e.g., sex and age), f) power calculation reported and the study was adequately powered to detect hypothesised relationships. Next, percentage agreement and Cohen's kappa coefficient⁴⁶ were calculated and discrepancies were discussed until 100% agreement was reached.

Evidence Synthesis

As shown in Figure 1, the literature search yielded 13,435 studies. Following title and abstract screening, 373 full-text articles were retrieved and reviewed. Ninety-eight studies met the inclusion criteria and at least one suitable effect size could be calculated for all but one study.⁴⁷

Study Characteristics

Study characteristics are presented in Table S2. Publication dates ranged from 1988-2015, with 66% being published in the last 5 years. Studies were mostly observational (98%), as experimental studies predominantly conduct interventions in laboratory settings rather than a specific life domain. The observational studies were mostly cross-sectional (76%), the sample sizes ranged from 19 to 106,319, with a total sample size of 648,726, and the majority of studies (76%) sampled adults. The majority of studies measured leisure-time PA (95%) – 13% measured work-related PA, 14% measured transport PA, 15% measured household PA,

3% measured school sport, and 3% measured physical education. Table S3 shows the number of studies measuring each mental health variable within each PA domain by age. Finally, all studies in this review exclusively utilised self-report measures of PA such as reporting PA participation over the previous days, weeks, or months.

Effect Sizes, Heterogeneity, and Moderator Analyses

Figure 2 highlights the effect size and 95% CI for each PA domain for mental health and mental ill-health and full results of the moderated meta-analyses are presented in Table 1 and Table 2.

Physical activity has a weak negative association with mental ill-health ($r = -.09$, 95% CI = $-.13, -.04$, $p < .001$). Overall, PA domain significantly moderates the strength of the relationship between PA and mental ill-health ($p < .001$) and explains 46% of the variance. In comparison to leisure-time PA, school sport is not significantly different ($\beta = -.04$, $p > .05$), whereas work-related PA ($\beta = .19$, $p < .001$), transport PA ($\beta = .07$, $p < .05$), household PA ($\beta = .11$, $p < .001$), and physical education ($\beta = .07$, $p < .05$) are all positive predictors of the effect size.

Leisure-time PA has a weak inverse association with mental ill-health and large heterogeneity is evident. Table 1 shows that the outcome measured is a significant predictor of strength, accounting for 36% of the variance. This result is due to a stronger negative association between leisure-time PA and depression ($r = -.16$), compared to other mental ill-health measures ($r = -.03$ to $-.09$). School sport is also significantly associated with mental ill-health, however, the effect ($r = -.09$) is slightly smaller than leisure-time ($r = -.11$). In contrast, work-

related PA has a weak positive association with mental ill-health and age is a significant predictor ($p = .03$) as the association is larger among adults ($r = .10$) than adolescents ($r = .01$), but this only explains 13% of the variance. Transport PA, household PA, and physical education have no significant association with mental ill-health.

Overall, PA in the included studies has a weak positive association with mental health ($r = .11$, 95% CI = .07 to .17, $p < .001$). Physical activity domain does not significantly moderate the relationship between PA and mental health ($p = .59$), as work-related PA ($\beta = -.03$), household PA ($\beta = -.03$), school sport ($\beta = -.03$), and physical education ($\beta = -.03$) are all non-significant ($p > .05$) negative predictor coefficients, compared to leisure-time. Nevertheless, the effect sizes varied between domains.

Leisure-time PA has a weak positive association with mental health and the effect size is not significantly moderated by the mental health outcome, sex, age, study design, PA measure, or risk of bias. Transport PA also has a weak positive association with mental health, but the CI is much wider when compared to leisure-time PA. The purpose of the trip (e.g., walking to work, walking during leisure-time) is the only significant moderator ($p = .01$) and explains majority of the variation in effect sizes ($R^2 = 76\%$). More specifically, active travel to and from work has a stronger relationship with mental health ($r = .28$) than active travel where the purpose was not identified or where all active trips were measured together ($r = .05$). While work-related PA is positively associated with mental health, heterogeneity is high and the relationship is not significant. Further, age is a significant moderator ($p = .002$) and explains 89% of the variance, as work-related PA is only positively associated with mental health in the one study examining adults >65 years ($r = .37$). Household PA, school sport, and physical education are not significantly associated with mental health.

Risk of Bias

Initial agreement between the two reviewers was 90% and the kappa coefficient ($\kappa = .81$) indicated strong agreement.⁴⁸ Upon discussion 100% agreement was achieved. Nineteen percent of studies have a high risk of bias and 59% have a moderate risk of bias, while only 18% have a low risk of bias. The complete results for risk of bias within studies are shown in Table S4. Non-significant Egger's test results for the mental health ($t = 1.6121$, $p = .11$) and mental ill-health meta-analyses ($t = 0.0925$, $p = .93$) revealed low asymmetry, suggesting low risk of bias across studies.

Discussion

The key message from this meta-analysis is that PA is not consistently associated with mental health or mental ill-health across different life domains. In fact, life domain significantly moderates the relationship between PA and mental ill-health. This is an important finding given the increased focus on lifestyle physical activity (e.g., standing desks and parking further from destinations to encourage active travel) in recent years and the existence of PA guidelines that encourage individuals to participate in PA in various domains.⁴⁹ While lifestyle PA outside leisure-time may improve people's physical health, such behaviours may not benefit mental health.⁵⁰ Promoting PA specifically during leisure-time, compared to other PA domains, may be most beneficial in terms of mental health promotion and the prevention of mental ill-health. However, it is also possible that individuals with poor mental health choose to participate in less leisure-time PA, whereas the same does not occur for other domains such as household PA. Age, sex, types of measurements, and study design explain little heterogeneity. As such, it is possible that other factors are causing some of the variation

between domains and within domains, and the results of the current review highlight the need to understand the source of this variation.

Evidence supports a number of psychosocial mechanisms that explain the effect of PA on mental health; however, there is little understanding as to whether these mechanisms play the same role in different PA domains.²² For example, PA enhances self-efficacy and exposes individuals to challenges that offer opportunities to develop confidence and a sense of mastery.^{51,52} Domains such as work-related PA and household PA may consist primarily of walking, and given the simplicity of that activity, PA in these domains appears likely to offer fewer opportunities to experience a sense of mastery than PA during leisure-time.⁵³

Alternatively, the social interaction hypothesis suggests that participating in PA with other people may influence mental health outcomes.⁵⁴ Transport PA may or may not be conducted with other people and so it is possible that social interaction explains some of the variance within this domain. The distraction hypothesis suggests that PA diverts individuals' attention from stressful life events which leads to improved mental wellbeing.⁵⁵ Perhaps, leisure-time PA provides a distraction from stress, but PA during domains such as work do not; indeed, work itself may be a source of stress.

Another potentially important factor is motivation. Leisure-time PA is often a chosen activity in which individuals are likely to participate due to enjoyment or perceived benefits.^{56,57}

Choosing to participate for enjoyment is central to autonomous motivation.⁵⁸ Alternatively, work-related PA is likely to be viewed as a compulsory task and may be enforced by another person or only carried out for the purpose of receiving an external reward (i.e., pay).¹³ If an activity is undertaken due to such factors, motivation is controlled.⁵⁸ Self-determination theory posits that behaviours which are undertaken due to autonomous motivation are more

likely to be associated with the satisfaction of psychological needs (i.e., autonomy, competence, and relatedness), and that wellbeing is enhanced when an individual's psychological needs are satisfied.⁵⁹ As such, autonomously motivated behaviours are expected to yield more positive psychological outcomes than behaviours undertaken due to controlled motivation.^{58,60,61} Further, evidence also shows that exercise involving choice is associated with increased positive affect and enjoyment, compared to exercise with no choice.⁶²⁻⁶⁴

Self-determined motivation may also explain some of the variation within each PA domain.¹⁷ While transport PA is positively associated with mental health among adults, no such relationship was found for adolescents. It is possible that this finding reflects the mental health benefits experienced when adults personally choose to walk to work because it is enjoyable or important, while young people who are forced to walk to school do not experience the same benefit. Furthermore, Asztalos reported that transport PA was associated with increased stress among blue collar workers but not among white collar workers.¹³ Financial factors may force some blue collar workers to bicycle to work (i.e., controlled motivation), while white collar workers may be more likely to own a car, yet autonomously choose to walk or ride due to health benefits or enjoyment.¹³ However, the role of autonomous motivation in the relationship between PA and mental health has not yet been explored.

Increasing PA during work is not likely to be worthwhile in terms of reducing the prevalence of mental ill-health, as individuals with occupations that involve higher amounts of PA, such as blue collar jobs, are more likely to experience mental ill-health. This finding suggests that PA is not automatically associated with greater mental health and reduced mental ill-health,

and that contextual factors are crucial to such relationships. It is possible that work-related PA does not provide a distraction from stress or provide opportunities for improved self-esteem, thereby not improving mental health. However, it is also possible that people with poorer mental health have limited occupational opportunities and secure employment in lower paid physically active jobs. Interestingly, work-related PA was positively associated with mental health as well as mental ill-health. While this result is plausible given that mental health and mental ill-health are not always negatively correlated,⁶⁵ it demonstrates that work-related PA is associated with mental health for some people and mental ill-health for others. This finding highlights the need for more extensive investigations of work-related PA in order to determine when it may be beneficial, as opposed to detrimental.

This review provides the first meta-analytic evidence of the relationships between PA undertaken in different life domains, and mental health and mental ill-health. Consequently, it provides the most comprehensive evidence showing that leisure-time PA is likely to be the optimal domain to promote mental health and prevent mental ill-health. The number of studies and, therefore, the total combined sample size, are both strengths of this study, as is the range of outcomes included. Nevertheless, some limitations should also be noted. Firstly, 98% of the included studies were observational, the majority of which were cross-sectional. As cross sectional studies cannot infer causality, the study designs of the included studies are a limitation.⁶⁶ As such, well designed experimental studies are needed to improve understanding of the causal relationships in different domains. Secondly, all 98 studies in this review utilised self-report measures of PA. This is a limitation of the current review and highlights the need for objective measures of domain-specific PA. While objectively measuring PA within specific domains is challenging and costly, advancements in technology have made it increasingly possible as Global Positioning Systems and wearable cameras have been used to measure PA within specific life domains.⁶⁷⁻⁷²

The small number of studies included for work-related PA, transport PA, household PA, school sport, and physical education, in comparison to leisure-time, is also a limitation as a number of moderator analyses within these domains could not be conducted due to insufficient studies. Additionally, Sallis' guideline for interpreting systematic review results states that evidence is inconclusive if less than four studies are included.⁷³ While school sport was inversely associated with mental ill-health, only two studies were included. Furthermore, while no relationship between physical education and mental health or mental ill-health was identified, there is not enough evidence at this stage to conclude that physical education is not associated with mental health. This lack of evidence warrants further investigation of these two domains. Additionally, heterogeneity remained high in majority of the sub-group meta-analyses suggesting that variation in the effect sizes was not consistently due to age, sex, study design, measures used, or risk of bias. Not identifying the source of this variation is a limitation of the current study. As such, other factors (e.g., motivation and social interaction) need to be explored. Finally, studies included in this review measured the relationship between habitual PA and mental health. As it is possible that PA in some domains has an immediate effect on mood following exercise, despite not providing longer-term benefits, future studies need to explore acute effects of domain-specific PA.

Conclusion

Despite the well accepted relationship between PA and mental health, this review concludes that the relationship varies between different PA domains. Gaining knowledge of the specific factors that mediate or moderate these relationships could lead to the development of contextually tailored interventions and PA guidelines, and improve the effectiveness of PA as a prevention and treatment method. Nevertheless, the results of this meta-analyses indicate

that, compared to other PA domains, promoting PA during leisure-time is likely to be the most effective method of preventing mental ill-health.

Acknowledgements

This research was funded by an Australian Research Council Discovery Grant (DP130104659) awarded to Chris Lonsdale and David Lubans. During the course of this research, Rhiannon White was supported by an Australian Postgraduate Award and an Australian Catholic University Postgraduate Award. Philip Parker was supported by an Australian Research Council Discovery Early Career Researcher Award. David Lubans was supported by an Australian Research Council Future Fellowship. Thomas Astell-Burt was supported by a Postdoctoral Fellowship from the National Heart Foundation of Australia .

RLW led the design of the study, completed all searches, screening, and data collection, performed the statistical analyses, and led the writing of the manuscript. MB completed all screening of studies, cross-checked the data extraction, and helped draft the manuscript. PP participated in the statistical analyses and helped to draft the manuscript. DL was involved in the design of the study and helped draft the manuscript. TAB was involved in the design of the study and reviewed the manuscript. CL participated in the design of the study, was involved in the statistical analyses, and helped to draft the manuscript. All authors read and approved the final manuscript.

Type of PA measure							.26	37%
Risk of bias							.65	11%
School sport Outcome	2	3	-.09***	-.11, -.07	61.37%	26	.12	0%
Study design							.85	0%
Type of PA measure							1.0	0%
Risk of bias							.85	0%
Physical education Outcome	2	3	-.02	-.11, .08	97.70%		.002	100%
Negative affect	1	2	-.08**	-.14, -.03	n/a	n/a		
Depression	1	1	.05	-.09, .19	n/a	.14		
Type of PA measure							.09	100%
Risk of bias							.002	100%
Total score /6							.72	

Note: Boldface indicates statistical significance (*p<0.05, **p<0.01, ***p<0.001.

β , standardised regression coefficient; ES, effect sizes; I^2 , I^2 statistic indicating proportion of variation due to heterogeneity; k , number of studies; PA, physical activity; r , correlation effect size; R^2 , proportion of heterogeneity that can be explained by the moderator variable

Table 2

Domain Specific PA and Mental Health Meta-Analyses and Moderator Analyses

	<i>k</i>	# ES	<i>r</i>	95% CI	<i>I</i> ²	Fail-safe <i>N</i>	Predictor coefficient β	ANOVA <i>p</i> value	<i>R</i> ²
Leisure-time PA	33	58	.13***	.08, .18	99.59%	1, 931			
Outcome								.46	16%
Age								.96	2%
Sex								.50	14%
Study design								.65	1%
Type of PA measure								.58	34%
Risk of bias								.94	
Work-related PA	5	6	.10	-.02, .22	89.13%				
Outcome								.10	67%
Age								.002	89%
Adults	3	3	.02	-.09, .12	76.50%		n/a		
Older adults	1	1	.37***	.17, .54	n/a		.36***		
Sex								.70	9%
Study design								.80	1%
Type of PA measure								.03	95%
Duration	1	1	-.03	-.12, .06	n/a		-.09		
Frequency	1	1	.40***	.26, .54	n/a		.34***		
MET-hr/week	4	4	.06*	.01, .11	75.38%		n/a		
Risk of bias								.32	20%
Transport-related PA	7	11	.13*	.02, .23	99.01%	37			
Outcome								.37	27%
Age								.44	21%
Sex								.70	10%
Purpose of trip								.01	76%
Not specified, or includes all travel	5	9	.05	-.02, .12	86.80%		n/a		
To work	2	2	.28***	.19, .38	70.28%		.24***		
Type of PA measure								.03	76%
Duration	1	1	.48***	.28, .68	n/a		.46***		
Frequency	1	1	.27*	.04, .50	n/a		.25		
MET-hr/week	3	4	.02	-.09, .13	90.71%		n/n		
Binary	2	5	.17*	.03, .30	99.53%		.15		
Risk of bias								.15	27%
Household PA	4	7	.04	-.02, .09	97.56%				
Outcome								.03	100%
Mental health	1	2	.00	.09, .17	n/a		n/a		

Mental wellbeing	2	4	.01	-.04, .06	98.62%	-.12**		
Life satisfaction	1	1	-.01	-.06, .08	n/a	-.14**		
Sex							.20	84%
Study design							.06	24%
Type of PA measure							.46	69%
Risk of bias							.25	5%
School sport	2	5	-.04	-.29, .23	99.78%			
Outcome							.36	15%
Sex							.13	55%
Study design							.36	15%
Physical education	2	10	.05	-.02, .13	96.27%			
Sex							1.0	0%
Type of PA measure							.77	2%

Note: Boldface indicates statistical significance (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).
 β , standardised regression coefficient; ES, effect sizes; I^2 , I^2 statistic indicating proportion of variation due to heterogeneity; k , number of studies; PA, physical activity; r , correlation effect size; R^2 , proportion of heterogeneity that can be explained by the moderator variable

Figure 1. Flow diagram of articles throughout the systematic review. PA = physical activity.

Figure 2. Correlation and 95% CI within specific PA domains for mental health and mental ill-health. PA = physical activity.

References

*References marked with an asterisk indicate studies included in the meta-analysis.

1. Prince M, Patel V, Saxena S, et al. Global mental health 1: No health without mental health. *Lancet*. 2007;370:859-877.
2. Whiteford HA, Degenhardt L, Rehm J, et al. Global burden of disease attributable to mental and substance use disorders: Findings from the Global Burden of Disease Study 2010. *The Lancet*. 2013;382(9904):1575-1586.
3. Begg S, Vos T, Barker B, Stevenson C, Stanley L, Lopez AD. The burden of disease and injury in Australia 2003. Canberra: Australian Institute of Health and Welfare.; 2007.
4. World Health Organization. The World Health Report: 2001: Mental Health: New Understanding, New Hope. Geneva, Switzerland: World Health Organization.; 2001.
5. Herrman H, Saxena S, Moodie R. *Promoting mental health: Concepts, emerging evidence, practice: A report of the World Health Organization, Department of Mental Health and Substance Abuse in collaboration with the Victorian Health Promotion Foundation and the University of Melbourne*. Geneva, Switzerland: World Health Organization; 2005.
6. Bize R, Johnson JA, Plotnikoff RC. Physical activity level and health-related quality of life in the general adult population: A systematic review. *Prev Med*. 2007;45(6):401-415.
7. Biddle SJH, Asare M. Physical activity and mental health in children and adolescents: A review of reviews. *Br J Sports Med*. 2011;45(11):886-895.
8. Janssen I, LeBlanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act*. 2010;7(40):1-16.
9. Shephard RJ. Limits to the measurement of habitual physical activity by questionnaires. *Br J Sports Med*. 2003;37:197-206.
10. Sallis JF, Cervero RB, Ascher W, Henderson KA, Kraft MK, Kerr J. An ecological approach to creating active living communities. *Annu Rev Public Health*. 2006;27:297-322.
11. Pratt M, Macera CA, Sallis JF, O'Donnell M, Frank LD. Economic interventions to promote physical activity: Application of the SLOTH model. *Am J Prev Med*. 2004;27(3):136-145.
12. Teychenne M, Ball K, Salmon J. Physical activity and likelihood of depression in adults: A review. *Prev Med*. 2008;46:397-411.
- *13. Asztalos M, Wijndaele K, De Bourdeaudhuij I, et al. Specific associations between types of physical activity and components of mental health. *J Sci Med Sport*. 2009;12(4):468-474.
- *14. Kull M, Ainsaar M, Kiive E, Raudsepp L. Relationship Between Low Depressiveness and Domain Specific Physical Activity in Women. *Health Care Woman Int*. 2012;33(5):457-472.
- *15. Ohta M, Mizoue T, Nishima N, Ikeda M. Effect of the physical activities in leisure time and commuting to work on mental health. *Journal of Occupational Health*. 2007;49:46-52.
16. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Ann Intern Med*. 2009;151(4):264-269.
17. Cattan M. *Mental Health And Well Being In Later Life*. McGraw-Hill International; 2009.

18. Luhmann M, Hawkey LC, Eid M, Cacioppo JT. Time frames and the distinction between affective and cognitive well-being. *J Res Pers.* 2012;46(4):431-441.
19. Barry MM, Jenkins R. *Implementing mental health promotion.* Philadelphia: Churchill Livingstone Elsevier; 2007.
20. Fontaine KL. *Mental health nursing.* 6 ed. Upper Saddle River: Pearson Prentice Hall; 2009.
21. World Health Organization. Promoting mental health: Concepts, emerging evidence, practice. Summary report. Geneva, Switzerland: World Health Organization.; 2004.
22. Lubans D, Richards J, Hillman C, et al. Physical activity for cognitive and mental health in youth: A systematic review of mechanisms. *Pediatrics.* 2016;138(3):e20161642.
23. Goldberg D, Krueger R, Andrews G, Hobbs M. Emotional disorders: cluster 4 of the proposed meta-structure for DSM-V and ICD-11. *Psychol Med.* 2009;39(12):2043-2059.
24. Tut TG, Wang JL, Lim CCL. Negative association between T102C polymorphism at the 5-HT2A receptor gene and bipolar affective disorders in Singaporean Chinese. *J Affect Disord.* 2000;58:211-214.
25. McLoone J, Hudson JL. Treating anxiety disorders in a school setting. *Educ Treat Children.* 2006;29(2):219-242.
26. Krueger RF, McGue M, Iacono WG. The higher-order structure of common DSM mental disorders: Internalization, externalization, and their connections to personality. *Pers Individ Dif.* 2001;30(7):1245-1259.
27. Borenstein M, Hedges LV, Higgins JPTH, Rothstein HR. *Introduction to meta-analysis.* West Sussex: John Wiley & Sons; 2009.
28. Rosenthal R. *Meta-analytic procedures for social research.* Newbury Park: Sage; 1991.
29. Hunter JE, Schmidt FL. *Methods of meta-analysis: Correcting error and bias in research findings.* 2nd ed. London: Sage Publications; 2004.
30. Charles EP. The correction for attenuation due to measurement error: clarifying concepts and creating confidence sets. *Psychol Methods.* 2005;10(2):206-226.
31. Cohen J, Cohen P, West SG, Aiken LS. *Applied multiple regression/correlation analysis for the behavioral sciences.* New Jersey: Routledge; 2013.
32. Cheung MWL. *Meta-analysis: A structural equation modeling approach.* West Sussex, United Kingdom: John Wiley & Sons; 2015.
33. Cheung MWL. metaSEM: An R package for meta-analysis using structural equation modeling. *Front Psychol.* 2015;5(1521).
34. Cohen J. *Statistical power analysis for the behavioral sciences.* New York: Academic Press; 1977.
35. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Br Med J.* 2003;327(7414):557-560.
36. Rücker G, Schwarzer G, Carpenter JR, Schumacher M. Undue reliance on I² in assessing heterogeneity may mislead. *BMC Med Res Methodol.* 2008;8(1):79-87.
37. Higgins JP, Green S. *Cochrane handbook for systematic reviews of interventions (Vol. 5).* Chichester: Wiley Online Library; 2008.
38. Doherty D, Kaufman M, Lindsay D, Panzer L, Sharpe D, Tonkin R. Age limits and adolescents [policy statement]. Ottawa: Canadian Paediatric Society.; 1994.
39. World Health Organization. Definition of an older or elderly person. Geneva, Switzerland: World Health Organization.; 2012.
40. Rosenthal R. The file drawer problem and tolerance for null results. *Psychol Bull.* 1979;86(3):638-641.

41. Sterne JA, Gavaghan D, Egger M. Publication and related bias in meta-analysis: power of statistical tests and prevalence in the literature. *J Clin Epidemiol.* 2000;53(11):1119-1129.
42. Von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Prev Med.* 2007;45(4):247-251.
43. Moher D, Schulz KF, Altman DG. The CONSORT statement: Revised recommendations for improving the quality of reports of parallel group randomized trials. *BMC Med Res Methodol.* 2001;1(1).
44. Babic MJ, Morgan PJ, Plotnikoff RC, Lonsdale C, White RL, Lubans DR. Physical activity and physical self-concept in youth: Systematic review and meta-analysis. *Sports Med.* 2014;44(11):1589-1601.
45. Owen KB, Parker PD, Van Zanden B, MacMillan F, Astell-Burt T, Lonsdale C. Physical activity and school engagement in youth: A systematic review and meta-analysis. *Educ Psychol.* 2016:1-17.
46. Cohen J. Weighted kappa: Nominal scale agreement provision for scaled disagreement or partial credit. *Psychol Bull.* 1968;70(4):213-220.
47. Bogaert I, De Martelaer K, Deforche B, Clarys P, Zinzen E. Associations between different types of physical activity and teachers' perceived mental, physical, and work-related health. *BMC Public Health.* 2014;14(1).
48. McHugh ML. Interrater reliability: The kappa statistic. *Biochemia Medica.* 2012;22(3):276-282.
49. Department of Health and Ageing. National Physical Activity Guidelines for Australians. Canberra: Commonwealth of Australia.; 2014.
50. Dunn AL. Effectiveness of lifestyle physical activity interventions to reduce cardiovascular disease. *Am J Lifestyle Med.* 2009;3(1 suppl):11S-18S.
51. Craft LL, Perna FM. The benefits of exercise for the clinically depressed. *Prim Care Companion J Clin Psychiatry.* 2004;6(3):104.
52. Paluska SA, Schwenk TL. Physical activity and mental health. *Sports Med.* 2000;29(3):167-180.
53. Sohn AJ, Hasnain M, Sinacore JM. Impact of exercise (walking) on blood pressure levels in African American adults with newly diagnosed hypertension. *Ethn Dis.* 2007;17(3):503-507.
54. Bailey M, McLaren S. Physical activity alone and with others as predictors of sense of belonging and mental health in retirees. *Aging and Mental Health.* 2005;9(1):82-90.
55. Leith LM. *Foundations of exercise and mental health.* Morgantown: Fitness Information Technology Morgantown, WV; 2010.
56. Seippel Ø. The meanings of sport: Fun, health, beauty or community? *Sport in Society.* 2006;9(1):51-70.
57. McCarthy PJ, Jones MV, Clark-Carter D. Understanding enjoyment in youth sport: A developmental perspective. *Psychol Sport Exerc.* 2008;9(2):142-156.
58. Ryan RM, Deci EL. Intrinsic and extrinsic motivations: Classic definitions and new directions. *Contemp Educ Psychol.* 2000;25(1):54-67.
59. Weinstein N, Ryan RM. When helping helps: autonomous motivation for prosocial behavior and its influence on well-being for the helper and recipient. *J Pers Soc Psychol.* 2010;98(2):222-244.
60. Deci EL, Ryan RM. Facilitating optimal motivation and psychological well-being across life's domains. *Can Psychol.* 2008;49(1):14-34.
61. Deci EL, Ryan RM. Self-determination theory: A macrotheory of human motivation, development, and health. *Can Psychol.* 2008;49(3):182-185.

62. Parfitt G, Rose EA, Burgess WM. The psychological and physiological responses of sedentary individuals to prescribed and preferred intensity exercise. *Br J Health Psychol.* 2006;11(1):39-53.
63. Vazou-Ekkekakakis S, Ekkekakis P. Affective consequences of imposing the intensity of physical activity: does the loss of perceived autonomy matter. *Hellenic Journal of Psychology.* 2009;6(2):125-144.
64. Lubans DR, Smith JJ, Morgan PJ, et al. Mediators of psychological well-being in adolescent boys. *J Adolesc Health.* 2016;58(2):230-236.
65. Keyes CL. Mental illness and/or mental health? Investigating axioms of the complete state model of health. *J Consult Clin Psychol.* 2005;73(3):539-548.
66. Kirk RE. *Experimental design: Procedures for the behavioral sciences.* 4th ed. Los Angeles: SAGE Publications; 2013.
67. Ridgers ND, Fairclough S. Assessing free-living physical activity using accelerometry: practical issues for researchers and practitioners. *Eur J Sport Sci.* 2011;11(3):205-213.
68. Schipperijn J, Andersen HB, Klinker CD, et al. Locating schoolyard physical activity—Using GPS, accelerometry and GIS. *J Sci Med Sport.* 2012;15:S255.
69. Cooper AR, Andersen LB, Wedderkopp N, Page AS, Froberg K. Physical activity levels of children who walk, cycle, or are driven to school. *Am J Prev Med.* 2005;29(3):179-184.
70. Duncan MJ, Badland HM, Mummery WK. Applying GPS to enhance understanding of transport-related physical activity. *J Sci Med Sport.* 2009;12:549-556.
71. Rodriguez DA, Cho G, Elder JP, et al. Identifying walking trips from GPS and accelerometer data in adolescent females. *J Phys Act Health.* 2012;9(3):421-431.
72. Ermes M, Parkka J, Mantyjarvi J, Korhonen I. Detection of daily activities and sports with wearable sensors in controlled and uncontrolled conditions. *Information Technology in Biomedicine.* 2008;12(1):20-26.
73. Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc.* 2000;32(5):963-975.
- *74. Aldana SG, Sutton LD, Jacobson BH, Quirk MG. Relationships between leisure time physical activity and perceived stress. *Percept Mot Skills.* 1996;82(1):315-321.
- *75. Armstrong K, Edwards H. The effectiveness of a pram-walking exercise programme in reducing depressive symptomatology for postnatal women. *Int J Nurs Pract.* Aug 2004;10(4):177-194.
- *76. Balboa-Castillo T, León-Muñoz LM, Graciani A, Rodriguez-Artalejo F, Guallar-Castillon P. Longitudinal association of physical activity and sedentary behavior during leisure time with health-related quality of life in community-dwelling older adults. *Health Qual Life Outcomes.* 2011;9:47.
- *77. Barrington WE, Ceballos RM, Bishop SK, McGregor BA, Beresford SAA. Perceived stress, behavior, and body mass index among adults participating in a worksite obesity prevention program, seattle, 2005-2007. *Prev Chronic Dis.* 2012;9(10).
- *78. Bernaards CM, Jans MP, Van Den Heuvel SG, Hendriksen IJ, Houtman IL, Bongers PM. Can strenuous leisure time physical activity prevent psychological complaints in a working population? *Occup Environ Med.* 2006;63(1):10-16.
- *79. Bertheussen GF, Romundstad PR, Landmark T, Kaasa S, Dale O, Helbostad JL. Associations between physical activity and physical and mental health-A HUNT 3 study. *Med Sci Sports Exerc.* 2011;43(7):1220-1228.
- *80. Binsinger C, Laure P, Ambard MF. Regular extra curricular sports practice does not prevent moderate or severe variations in self-esteem or trait anxiety in early adolescents. *Journal of Sports Science and Medicine.* 2006;5(1):123-129.

- *81. Birkeland MS, Torsheim T, Wold B. A longitudinal study of the relationship between leisure-time physical activity and depressed mood among adolescents. *Psychol Sport Exerc.* 2009;10(1):25-34.
- *82. Borges LJ, Benedetti TRB, Xavier AJ, deOrsi E. Associated factors of depressive symptoms in the elderly: EpiFloripa study. *Rev Saude Publica.* 2013;47(4).
- *83. Branco JC, Jansen K, Oses JP, et al. Practice of leisure-time physical activities and episodes of mood alteration amongst men and women. *J Affect Disord.* 2014;169:165-169.
- *84. Brooks FM, Smeeton NC, Chester K, Spencer N, Klemmera E. Associations between physical activity in adolescence and health behaviours, well-being, family and social relations. *Int J Health Promot Educ.* 2014;52(5):271-282.
- *85. Brunet A, Augestad LB, Gudmundsdottir SL. Personality, physical activity, and symptoms of anxiety and depression: The HUNT study. *Soc Psychiatry Psychiatr Epidemiol.* 2013;48(5):745-756.
- *86. Brunet A, Gudmundsdottir SL, Augestad LB. Gender-specific associations between leisure-time physical activity and symptoms of anxiety: The HUNT study. *Soc Psychiatry Psychiatr Epidemiol.* 2014;50(3):419-427.
- *87. Bültmann U, Kant I, Kasl SV, Schröer KAP, Swaen GMH, Van den Brandt PA. Lifestyle factors as risk factors for fatigue and psychological distress in the working population: Prospective results from the Maastricht cohort study. *J Occup Environ Med.* 2002;44(2):116-124.
- *88. Bustamante EE, Wilbur J, Marquez DX, Fogg L, Staffileno BA, Manning A. Physical activity characteristics and depressive symptoms in older Latino adults. *Ment Health Phys Act.* 2013;6(2):69-77.
- *89. Buttery AK, Mensink GB, Busch MA. Healthy behaviours and mental health: findings from the German Health Update (GEDA). *Eur J Public Health.* Jul 23 2014.
- *90. Cerin E, Leslie E, Sugiyama T, Owen N. Associations of multiple physical activity domains with mental well-being. *Ment Health Phys Act.* 2009;2(2):55-64.
- *91. Chen LJ, Stevinson C, Ku PW, Chang YK, Chu DC. Relationships of leisure-time and non-leisure-time physical activity with depressive symptoms: A population-based study of Taiwanese older adults. *Int J Behav Nutr Phys Act.* 2012;9.
- *92. Chi SH, Wang JY, Tsai AC. Combined association of leisure-time physical activity and fruit and vegetable consumption with depressive symptoms in older Taiwanese: Results of a national cohort study. *Geriatr Gerontol Int.* 2015;16(2):244-251.
- *93. Choi J, Wilbur J, Miller A, Szalacha L, McAuley E. Correlates of leisure-time physical activity in Korean immigrant women. *West J Nurs Res.* 2008;30(5):620-638.
- *94. Craike MJ, Coleman D, MacMahon C. Direct and buffering effects of physical activity on stress-related depression in mothers of infants. *J Sport Exerc Psychol.* 2010;32(1):23-38.
- *95. Desha LN, Ziviani JM, Nicholson JM, Martin G, Darnell RE. Physical activity and depressive symptoms in American adolescents. *Journal of Sport and Exercise Psychology.* 2007;29(4):534-543.
- *96. Doerksen SE, Elavsky S, Rebar AL, Conroy DE. Weekly fluctuations in college student leisure activities and well-being. *Leisure Sciences.* 2014;36(1):14-34.
- *97. Dupuis SL, Smale BJA. An examination of relationship between psychological well-being and depression and leisure activity participation among older adults. *Society & Leisure / Loisir & Société.* 1995;18(1):67-92.
- *98. Eime R, Harvey J, Payne W. Dose-response of women's health-related quality of life (HRQOL) and life satisfaction to physical activity. *Journal of Physical Activity and Health.* 2014;11(2):330-338.

- *99. Feuerhahn N, Sonnentag S, Woll A. Exercise after work, psychological mediators, and affect: A day-level study. *European Journal of Work and Organizational Psychology*. 2014;23(1):62-79.
- *100. Fisher KJ, Li F. A community-based walking trial to improve neighborhood quality of life in older adults: a multilevel analysis. *Ann Behav Med*. Dec 2004;28(3):186-194.
- *101. Fløtnes IS, Nilsen TIL, Augestad LB. Norwegian adolescents, physical activity and mental health: The Young-HUNT study. *Norsk Epidemiologi*. 2011;20(2):153-161.
- *102. Giacobbi Jr PR, Tuccitto DE, Frye N. Exercise, affect, and university students' appraisals of academic events prior to the final examination period. *Psychol Sport Exerc*. 2007;8(2):261-274.
- *103. Goldfield GS, Henderson K, Buchholz A, Obeid N, Hien N, Flament MF. Physical activity and psychological adjustment in adolescents. *J Phys Act Health*. 2011;8(2):157-163.
- *104. Griffiths LJ, Dowda M, Dezateux C, Pate R. Associations between sport and screen-entertainment with mental health problems in 5-year-old children. *Int J Behav Nutr Phys Act*. 2010;7.
- *105. Hamer M, Molloy GJ, de Oliveira C, Demakakos P. Leisure time physical activity, risk of depressive symptoms, and inflammatory mediators: The English Longitudinal Study of Ageing. *Psychoneuroendocrinology*. 2009;34(7):1050-1055.
- *106. Hamer M, Stamatakis E, Steptoe A. Dose-response relationship between physical activity and mental health: The Scottish Health Survey. *Br J Sports Med*. 2009;43(14):1111-1114.
- *107. Herman KM, Hopman WM, Sabiston CM. Physical activity, screen time and self-rated health and mental health in Canadian adolescents. *Prev Med*. 2015;73:112-116.
- *108. Humphreys DK, Goodman A, Ogilvie D. Associations between active commuting and physical and mental wellbeing. *Prev Med*. 2013;57(2):135-139.
- *109. Hyde AL, Conroy DE, Pincus AL, Ram N. Unpacking the feel-good effect of free-time physical activity: Between- and within-person associations with pleasant-activated feeling states. *Journal of Sport and Exercise Psychology*. 2011;33(6):884-902.
- *110. Im EO, Ham OK, Chee E, Chee W. Physical Activity and Depressive Symptoms in Four Ethnic Groups of Midlife Women. *West J Nurs Res*. May 29 2014.
- *111. Inal S, Subasi F, Ay SM, Hayran O. The links between health-related behaviors and life satisfaction in elderly individuals who prefer institutional living. *BMC Health Serv Res*. 2007;7(30).
- *112. Iwasaki Y, Zuzanek J, Mannell RC. The effects of physically active leisure on stress-health relationships. *Canadian Journal of Public Health*. 2001;92(3):214-218.
- *113. Jewett R, Sabiston CM, Brunet J, O'Loughlin EK, Scarapicchia T, O'Loughlin J. School sport participation during adolescence and mental health in early adulthood. *J Adolesc Health*. 2014;55(5):640-644.
- *114. Joseph RP, Royse KE, Benitez TJ, Pekmezi DW. Physical activity and quality of life among university students: Exploring self-efficacy, self-esteem, and affect as potential mediators. *Qual Life Res*. 2014;23(2):661-669.
- *115. Jurakić D, Pedišić Z, Greblo Z. Physical activity in different domains and health-related quality of life: A population-based study. *Qual Life Res*. 2010;19(9):1303-1309.
- *116. Kilpatrick M, Sanderson K, Blizzard L, Teale B, Venn A. Cross-sectional associations between sitting at work and psychological distress: Reducing sitting time may benefit mental health. *Ment Health Phys Act*. 2013;6(2):103-109.
- *117. Kim J, Chun S, Heo J, Lee S, Han A. Contribution of leisure-time physical activity on psychological benefits among elderly immigrants. *Appl Res Qual Life*. 2014;11:461-470.

- *118. Korniloff K, Vanhala M, Kautiainen H, et al. Lifetime leisure-time physical activity and the risk of depressive symptoms at the ages of 65-74 years: the FIN-D2D survey. *Prev Med.* May 2012;54(5):313-315.
- *119. Kremer P, Elshaug C, Leslie E, Toumbourou JW, Patton GC, Williams J. Physical activity, leisure-time screen use and depression among children and young adolescents. *J Sci Med Sport.* 2014;17(2):183-187.
- *120. Ku PW, Fox KR, Chen LJ. Physical activity and depressive symptoms in Taiwanese older adults: A seven-year follow-up study. *Prev Med.* 2009;48(3):250-255.
- *121. Kwag K, Martin P, Russell D, Franke W, Kohut M. The impact of perceived stress, social support, and home-based physical activity on mental health among older adults. *Int J Aging Hum Dev.* 2011;72(2):137-154.
- *122. Lee PL, Lan W, Lee CCL. Physical activity related to depression and predicted mortality risk: Results from the Americans' Changing Lives Study. *Educ Gerontol.* 2012;38(10):678-690.
- *123. Lin L, Halgin RP, Well AD, Ockene I. The relationship between depression and occupational, household, and leisure-time physical activity. *J Clin Sport Psychol.* 2008;2(2):95-107.
- *124. Lutz RS, Lochbaum MR, Lanning B, Stinson LG, Brewer R. Cross-lagged relationships among leisure-time exercise and perceived stress in blue-collar workers. *Journal of Sport and Exercise Psychology.* 2007;29(6):687-705.
- *125. Malabo A, Van Eeden C, Wissing MP. Sport participation, psychological wellbeing, and psychosocial development in a group of young black adults. *South African Journal of Psychology.* 2007;37(1):188-206.
- *126. Martin A, Goryakin Y, Suhrcke M. Does active commuting improve psychological wellbeing? Longitudinal evidence from eighteen waves of the British Household Panel Survey. *Prev Med.* 2014;69:296-303.
- *127. Martins LC, Lopes CS. Rank, job stress, psychological distress and physical activity among military personnel. *BMC Public Health.* 2013;13:716.
- *128. McKercher C, Patton GC, Schmidt MD, Venn AJ, Dwyer T, Sanderson K. Physical activity and depression symptom profiles in young men and women with major depression. *Psychosom Med.* 2013;75(4):366-374.
- *129. McKercher C, Schmidt MD, Sanderson K, Dwyer T, Venn AJ. Physical activity and depressed mood in primary and secondary school-children. *Ment Health Phys Act.* 2012;5(1):50-56.
- *130. McKercher CM, Schmidt MD, Sanderson KA, Patton GC, Dwyer T, Venn AJ. Physical Activity and Depression in Young Adults. *Am J Prev Med.* 2009;36(2):161-164.
- *131. Molarius A, Berglund K, Eriksson C, et al. Mental health symptoms in relation to socio-economic conditions and lifestyle factors--a population-based study in Sweden. *BMC Public Health.* 2009;9:302.
- *132. Moore KA, Babyak MA, Wood CE, et al. The association between physical activity and depression in older depressed adults. *Journal of Aging and Physical Activity.* 1999;7(1):55-61.
- *133. Motl RW, Birnbaum AS, Kubik MY, Dishman RK. Naturally occurring changes in physical activity are inversely related to depressive symptoms during early adolescence. *Psychosom Med.* 2004;66(3):336-342.
- *134. Mouchacca J, Abbott GR, Ball K. Associations between psychological stress, eating, physical activity, sedentary behaviours and body weight among women: A longitudinal study. *BMC Public Health.* 2013;13(1).
- *135. Mouton CP, Calmbach WL, Dhanda R, Espino DV, Hazuda H. Barriers and benefits to leisure-time physical activity among older Mexican Americans. *Arch Fam Med.* 2000;9(9):892-897.

- *136. Mutrie N, Hannah MK. The importance of both setting and intensity of physical activity in relation to non-clinical anxiety and depression. *Int J Health Promot Educ*. 2007;45(1):24-32.
- *137. Nakamura PM, Teixeira IP, Smirmaul BPC, et al. Health related quality of life is differently associated with leisure-time physical activity intensities according to gender: A cross-sectional approach. *Health and Quality of Life Outcomes*. 2014;12(1):1-10.
- *138. Pagán R. How do leisure activities impact on life satisfaction? Evidence for German people with disabilities. *Appl Res Qual Life*. 2014;10:557-572.
- *139. Pasco JA, Williams LJ, Jacka FN, et al. Habitual physical activity and the risk for depressive and anxiety disorders among older men and women. *Int Psychogeriatr*. 2011;23(2):292-298.
- *140. Pastor Y, Balaguer I, Pons D, Garcia-Merita M. Testing direct and indirect effects of sports participation on perceived health in Spanish adolescents between 15 and 18 years of age. *J Adolesc*. Dec 2003;26(6):717-730.
- *141. Pedišić Z, Greblo Z, Phongsavan P, Milton K, Bauman AE. Are total, intensity- and domain-specific physical activity levels associated with life satisfaction among university students? *PLoS One*. 2015;10(2):e0118137.
- *142. Peeters G, Van Gellecum YR, Van Uffelen JGZ, Burton NW, Brown WJ. Contribution of house and garden work to the association between physical activity and well-being in young, mid-aged and older women. *Br J Sports Med*. 2014;48(12):996-1001.
- *143. Petronyte G, Zaborskis A. The influence of individual and regional factors on association between leisure time physical activity and psychological complaints among adolescents in Europe. *Cent Eur J Public Health*. 2009;17(4):215-219.
- *144. Pickett K, Yardley L, Kendrick T. Physical activity and depression: A multiple mediation analysis. *Ment Health Phys Act*. 2012;5(2):125-134.
- *145. Purakom A, Nakornkhet K, Tanomsuk M, Pupanead S, André S, Joana C. Associations between physical activity, functional fitness, and mental health among older adults in Nakornpathom, Thailand. *Asian Journal of Exercise & Sports Science*. 2014;11(2):25-35.
- *146. Rocha SV, de Araújo TM, de Almeida MMG, Virtuoso Jr JS. Practice of physical activity during leisure time and common mental disorders among residents of a municipality of Northeast Brazil. *Revista Brasileira de Epidemiologia*. 2012;15(4):871-883.
- *147. Sabiston CM, O'Loughlin E, Brunet J, et al. Linking depression symptom trajectories in adolescence to physical activity and team sports participation in young adults. *Prev Med*. 2013;56(2):95-98.
- *148. Sanchez-Villegas A, Ara I, Dierssen T, Dela Fuente C, Ruano C, Martínez-González MA. Physical activity during leisure time and quality of life in a Spanish cohort: SUN (Seguimiento Universidad de Navarra) Project. *Br J Sports Med*. 2012;46(6):443-448.
- *149. Sanchez-Villegas A, Ara I, Guillén-Grima F, Bes-Rastrollo M, Varo-Cenarruzabeitia JJ, Martínez-González MA. Physical activity, sedentary index, and mental disorders in the sun cohort study. *Med Sci Sports Exerc*. 2008;40(5):827-834.
- *150. Sasidharan V, Payne L, Orsega-Smith E, Godbey G. Older adults' physical activity participation and perceptions of wellbeing: Examining the role of social support for leisure. *Managing Leisure*. 2006;11(3):164-185.
- *151. Schnohr P, Kristensen TS, Prescott E, Scharling H. Stress and life dissatisfaction are inversely associated with jogging and other types of physical activity in leisure time—The Copenhagen City Heart Study. *Scand J Med Sci Sports*. 2005;15(2):107-112.
- *152. Serrano-Sanchez JA, Lera-Navarro A, Espino-Torón L. Physical activity and differences of functional fitness and quality of life in older males. *Revista*

Internacional de Medicina y Ciencias de la Actividad Fisica y del Deporte.
2013;13(49):87-105.

- *153. Sieverdes JC, Ray BM, Sui X, et al. Association between leisure time physical activity and depressive symptoms in men. *Med Sci Sports Exerc.* 2012;44(2):260-265.
- *154. Stephens T. Physical activity and mental health in the United States and Canada: evidence from four population surveys. *Prev Med.* Jan 1988;17(1):35-47.
- *155. Surkan PJ, Ryan LM, Bidwell HW, Brooks DR, Peterson KE, Gillman MW. Psychosocial correlates of leisure-time physical activity in urban working-class adults. *J Phys Act Health.* 2005;2(4):397-411.
- *156. Tessier S, Vuillemin A, Bertrais S, et al. Association between leisure-time physical activity and health-related quality of life changes over time. *Prev Med.* 2007;44(3):202-208.
- *157. Teychenne M, Ball K, Salmon J. Associations between physical activity and depressive symptoms in women. *Int J Behav Nutr Phys Act.* 2008;5.
- *158. Teychenne M, Ball K, Salmon J. Physical activity, sedentary behavior and depression among disadvantaged women. *Health Educ Res.* 2010;25(4):632-644.
- *159. Teychenne M, Torres S, McNaughton SA, Salmon J, Ball K. Mediators of the relationship between sedentary behavior and depressive symptoms amongst disadvantaged women. *Ment Health Phys Act.* 2014;7(1):30-36.
- *160. Thøgersen-Ntoumani C, Fox KR, Ntoumanis N. Relationships between exercise and three components of mental well-being in corporate employees. *Psychol Sport Exerc.* 2005;6(6):609-627.
- *161. Valois RF, Zullig KJ, Huebner ES, Drane JW. Physical activity behaviors and perceived life satisfaction among public high school adolescents. *J Sch Health.* Feb 2004;74(2):59-65.
- *162. Vuillemin A, Boini S, Bertrais S, et al. Leisure time physical activity and health-related quality of life. *Prev Med.* 2005;41(2):562-569.
- *163. Wang F, DesMeules M, Luo W, Dai S, Lagace C, Morrison H. Leisure-Time Physical Activity and Marital Status in Relation to Depression Between Men and Women: A Prospective Study. *Health Psychol.* 2011;30(2):204-211.
- *164. Wendel-Vos GCW, Schuit AJ, Tijhuis MAR, Kromhout D. Leisure time physical activity and health-related quality of life: Cross-sectional and longitudinal associations. *Qual Life Res.* 2004;13(3):667-677.
- *165. Wijndaele K, Matton L, Duvinneaud N, et al. Association between leisure time physical activity and stress, social support and coping: A cluster-analytical approach. *Psychol Sport Exerc.* 2007;8(4):425-440.
- *166. Yang X, Telama R, Hirvensalo M, et al. Moderating effects of leisure-time physical activity on the association between job strain and depressive symptoms: The cardiovascular risk in young finns study. *J Occup Environ Med.* 2012;54(3):303-309.
- *167. Zullig KJ, White RJ. Physical activity, life satisfaction, and self-rated health of middle school students. *Appl Res Qual Life.* 2011;6(3):277-289.

Supplementary Text and Tables

Table S1.

Full list of search terms.

Physical Activity Keywords	Mental Health Keywords	Domain Specific Keywords		Excluded Terms		
“physical activit*”	“mental health”	domain*	domestic	Femur	Arousal	Ankle
“physical inactivity”	“mental well*being”	“types of”	housework	Patella	“heat stress”	Resistance
exercise	“psychological well*being”	“leisure time”	household	Knee	Ventricular	Rheumatology
sport*	“subjective well*being”	“leisure-time”	PE	Hip	“stress test*”	Lumbar
walking	“life satisfaction”	Sport*	P.E.	Lung	Renal	Gait
	“positive affect”	travel	“physical education”	Lactate	Heart	Rats
	“negative affect”	commut*	Occupation*	Contraction	Injur*	Metaboli*
	“mental illness”	transport*		Serum	Cardiovascular	Sclerosis
	“mental disorder*”	Transport-related		“Cartilage stress”	Diabetes	Stroke
	depress*	cycling		“Oxidative stress”	Trauma*	Dementia
	anxiety	walking		Cancer	Muscle	Biomechanics
	stress	work		Performance	Spin*	Ergonomics
	“psychological distress”	work-related		Horses	Arthritis	
				“Stress urinary incontinence”		

Full search extracted from Scopus

```

(( TITLE-ABS-KEY ( commut* ) OR TITLE-ABS-KEY ( "leisure time" ) OR TITLE-
ABS-KEY ( "leisure-time" ) OR TITLE-ABS-KEY ( "types of" ) OR TITLE-ABS-
KEY ( domain* ) OR TITLE-ABS-KEY ( work ) OR TITLE-ABS-
KEY ( occupation* ) OR TITLE-ABS-KEY ( travel ) OR TITLE-ABS-
KEY ( housework ) OR TITLE-ABS-KEY ( sport* ) OR TITLE-ABS-
KEY ( transport* ) OR TITLE-ABS-KEY ( cycling ) OR TITLE-ABS-
KEY ( domestic ) OR TITLE-ABS-KEY ( work-related ) OR TITLE-ABS-KEY ( transport-
related ) OR TITLE-ABS-KEY ( pe ) OR TITLE-ABS-KEY ( p.e. ) OR TITLE-ABS-
KEY ( "physical education" ) OR TITLE-ABS-
KEY ( household ) ) AND SUBJAREA ( mult OR medi OR nurs OR vete OR dent OR
heal OR mult OR arts OR busi OR deci OR econ OR psyc OR soci ) ) AND ( ( TITL
E-ABS-KEY ( "mental health" ) OR TITLE-ABS-KEY ( "mental well*being" ) OR TITLE-
ABS-KEY ( "psychological well*being" ) OR TITLE-ABS-KEY ( "subjective
well*being" ) OR TITLE-ABS-KEY ( "life satisfaction" ) OR TITLE-ABS-KEY ( "positive
affect" ) OR TITLE-ABS-KEY ( "negative affect" ) OR TITLE-ABS-KEY ( "mental
illness" ) OR TITLE-ABS-KEY ( "mental disorder*" ) OR TITLE-ABS-
KEY ( depress* ) OR TITLE-ABS-KEY ( anxiety ) OR TITLE-ABS-
KEY ( stress ) OR TITLE-ABS-KEY ( "psychological
distress" ) ) AND SUBJAREA ( mult OR medi OR nurs OR vete OR dent OR heal O
R mult OR arts OR busi OR deci OR econ OR psyc OR soci ) ) AND ( ( TITLE-ABS-
KEY ( "physical activit*" ) OR TITLE-ABS-KEY ( "physical inactivity" ) OR TITLE-
ABS-KEY ( exercise ) OR TITLE-ABS-KEY ( sport* ) OR TITLE-ABS-
KEY ( walking ) ) AND SUBJAREA ( mult OR medi OR nurs OR vete OR dent OR he
al OR mult OR arts OR busi OR deci OR econ OR psyc OR soci ) ) AND
NOT TITLE-ABS-KEY ( femur ) AND NOT TITLE-ABS-KEY ( patella ) AND
NOT TITLE-ABS-KEY ( knee ) AND NOT TITLE-ABS-KEY ( hip ) AND NOT TITLE-
ABS-KEY ( lung ) AND NOT TITLE-ABS-KEY ( lactate ) AND NOT TITLE-ABS-
KEY ( contraction ) AND NOT TITLE-ABS-KEY ( serum ) AND NOT TITLE-ABS-
KEY ( performance ) AND NOT TITLE-ABS-KEY ( "heat stress" ) AND NOT TITLE-
ABS-KEY ( arousal ) AND NOT TITLE-ABS-KEY ( "stress urinary incontinence" ) AND
NOT TITLE-ABS-KEY ( "cartilage stress" ) AND NOT TITLE-ABS-KEY ( renal ) AND
NOT TITLE-ABS-KEY ( "oxidative stress" ) AND NOT TITLE-ABS-KEY ( "stress
test*" ) AND NOT TITLE-ABS-KEY ( ventricular ) AND NOT TITLE-ABS-
KEY ( heart ) AND NOT TITLE-ABS-KEY ( injur* ) AND NOT TITLE-ABS-
KEY ( diabetes ) AND NOT TITLE-ABS-KEY ( muscle ) AND NOT TITLE-ABS-
KEY ( spine ) AND NOT TITLE-ABS-KEY ( arthritis ) AND NOT TITLE-ABS-
KEY ( trauma* ) AND NOT TITLE-ABS-KEY ( ankle ) AND NOT TITLE-ABS-
KEY ( sclerosis ) AND NOT TITLE-ABS-KEY ( stroke ) AND NOT TITLE-ABS-
KEY ( metaboli* ) AND NOT TITLE-ABS-KEY ( dementia ) AND NOT TITLE-ABS-
KEY ( biomechanics ) AND NOT TITLE-ABS-KEY ( ergonomics ) AND NOT TITLE-ABS-
KEY ( resistance ) AND NOT TITLE-ABS-KEY ( rheumatology ) AND NOT TITLE-ABS-
KEY ( lumbar ) AND NOT TITLE-ABS-KEY ( gait ) AND NOT TITLE-ABS-KEY )

```

Table S2.

Summary of studies included in the systematic review.

Author	Study Design	Country	Sample	Age (years)	Physical Activity Domain	Physical Activity Measure	Mental Health Measure
Aldana (1996)	Cross-sectional	United States	32,229 adults (57% female)	M = 37.3	Leisure time	1985 National Health Interview Survey	Strain Questionnaire
Armstrong (2004)	Experimental	Australia	19 adults (100% female)	Mo. = 30	Leisure time	n/a	Edinburgh Postnatal Depression Scale
Asztalos (2009)	Cross-sectional	Belgium	1,919 adults (47% female)	R = 20 - 65	Leisure time, Transport, Household	Flemish Physical Activity Computerized Questionnaire	GHQ12, Perceived Stress Scale
Balboa-Castillo, (2011)	Longitudinal	Spain	1,097 older aged (52% female)	M = 70.3 ± 5.6	Leisure time	Nurses' Health Study and the Health Professionals' Follow-up Study.	Spanish version of the SF-36
Barrington (2012)	Cross-sectional	United States	621 adults (57% female)	R = 18 - 65	Leisure time	Two items from the Godin Leisure-Time Exercise Questionnaire, one item from the IPAQ	Perceived Stress Scale-10
Bernaards (2006)	Longitudinal	Netherlands	1,747 adults (sex not reported)	Not reported	Leisure time	Frequency of PA	Dutch Version of the CES-D
Bertheussen (2011)	Cross-sectional	Norway	4,500 middle and older aged adults (56% female)	M = 53.1 ± 15	Leisure time	The number of times the participant exercised last week.	SF-8 Health Survey
Binsinger (2006)	Longitudinal	France	1,791 adolescents (53% female)	M = 11.1 ± 0.5	Leisure time	Do you do an extra curricular sport? Hours per week?	State-Trait Anxiety Inventory
Birkeland (2009)	Longitudinal	Norway	924 adolescents (45% female)	R = 13 - 23	Leisure time	One item	Seven item inventory
Bogaert (2014)	Cross-sectional	Belgium	1,066 adults (68% female)	M = 40.3 ± 9.7	Leisure time, Work, Transport, Household	IPAQ	SF-36
Borges (2013)	Cross-sectional	Brazil	1,656 older aged (64% female)	R = 60+	Leisure time	IPAQ	Geriatric Depression Scale
Branco (2014)	Cross-sectional	Brazil	1,953 young adults (55% female)	R = 18 - 35	Leisure time	IPAQ	Mini International Neuropsychiatric Interview 5.0 (MINI)
Brooks (2014)	Cross-sectional	England	4,404 adolescents (sex not reported)	Year 7, 9, and 11	Leisure time	Frequency and duration of vigorous exercise undertaken during leisure-time.	From the Health Behaviour in School-Aged Children World Health Organisation study
Brunes (2013)	Cross-sectional	Norway	38,743 adults (56% female)	M = 51.2 (females) M = 55.1 (males)	Leisure time	Frequency, duration, and intensity of weekly leisure-time PA over last 12 months. Self-report job rated as sedentary, frequent walking, frequent lifting, or heavy manual labor.	Hospital Anxiety and Depression Scale
Brunes (2014)	Longitudinal	Norway	23,991 adults (53% female)	M = 51.2 (females) M = 55.1 (males)	Leisure time	Frequency, duration, and intensity of weekly leisure-time PA over last 12 months.	Hospital Anxiety and Depression Scale – Anxiety Subscale

Bültmann (2002)	Longitudinal	Netherlands	7,208 adults (25% female)	R = 18 - 65	Leisure time	One item	GHQ12
Bustamante (2013)	Cross-sectional	United States	174 middle and older aged adults (74% female)	M = 65.97 ± 9.12	Leisure time, Household	40-item Community Healthy Activities Model Program for Seniors (CHAMPS) PA questionnaire	CES-D
Buttery (2014)	Cross-sectional	Germany	39,001 adults (56% female)	M = 48.1	Leisure time	One item	Health Related Quality of Life - 4, Self-reported depression diagnosis SF-12
Cerin (2009)	Longitudinal	Australia	2,194 adults (64% female)	M = 46.3 ± 12.0	Leisure time, Work, Transport, Household	IPAQ	
Chen (2012)	Cross-sectional	Taiwan	2,727 older aged (51% female)	R = 65+	Leisure time	Frequency and duration of sport.	10-item Chinese version of the CES-D
Chi (2015)	Longitudinal	Taiwan	2,630 middle and older aged adults (53% female)	R = 53+	Leisure time	Frequency, duration, and intensity of Leisure Time Physical Activity	CES-D
Choi (2008)	Cross-sectional	United States	188 adults (100% female)	M = 41 ± 10.96	Leisure time	IPAQ-Long Version	20-item Korean version of the CES-D
Craike (2010)	Cross-sectional	Australia	4,702 adults (100% female)	M = 34.6 ± 5.23	Leisure time	One item	"How difficult do you feel your life is at the moment?", Kessler Psychological Distress Scale
Desha (2007)	Cross-sectional	United States	727 adolescents (51% female)	M = 15.26	Leisure time	The Sport Involvement Index	Hovac's Children's Depression Inventory
Doerksen (2014)	Longitudinal	United States	33 adults (42% female)	R = 15+	Leisure time	Leisure Activity Scale	Satisfaction with Life Scale, Perceived Stress Scale, CES-D, Kuppens Affect Scale, State-Trait Anxiety Inventory
Dupuis (1995)	Cross-sectional	Canada	743 middle and older aged adults (58% female)	R = 55+	Leisure time	Frequency, duration, and intensity of participation in physically active leisure activities.	CES-D Bradburn Affect Scale
Eime (2014)	Cross-sectional	Australia	793 adults (100% female)	R = 18+	Leisure time	Duration of PA during previous week.	SF-36 , Life Satisfaction Score
Feuerhahn (2014)	Cross-sectional	Germany	126 adults (68% female)	M = 3778 ± 10.24	Leisure time	Exercise Diary	PANAS
Fisher (2004)	Experimental	United States	582 older aged adults (59% female)	R = 65+	Leisure time	3 items about neighborhood activity level.	SF-12
Fløtnes (2011)	Longitudinal	Norway	2,000 adolescents (52% female)	R = 13 - 19	Leisure time	Frequency, duration, and intensity of training, and type of activity.	Hopkins Symptom Check List-5 (SCL-5)
Giacobbi Jr (2007)	Cross-sectional	United States	59 young adults (37% female)	M = 21.56 ± 1.94	Leisure time	Leisure-time Exercise Questionnaire (LTEQ)	PANAS
Goldfield (2011)	Cross-sectional	United States	1,259 adolescents (59% female)	M = 14.8 ± 1.8	Leisure time	The Godin Leisure-Time Exercise Questionnaire	CDI, Multidimensional Anxiety Scale for Children (MASC-10)
Griffiths (2010)	Cross-sectional	United Kingdom	13,470 children (49% female)	M = 5.2	Leisure time	One item	Strengths and Difficulties Questionnaire (SDQ)
Hamer (2009)	Longitudinal	England	4,323 middle and older aged adults (52% female)	M = 63.4 ± 9.7	Leisure time	Frequency and intensity of physical activities during leisure time.	CES-D

Hamer (2009)	Cross-sectional	Scotland	19,842 adults (54% female)	$M = 45.2 \pm 15.5$	Leisure time, Household	Frequency of participation in ≥ 20 mins of leisure-time PA. Frequency and duration of household PA.	GHQ12
Herman (2015)	Cross-sectional	Canada	7,725 adolescents (49% female)	$R = 12 - 17$	Leisure time	Adaptation of Minnesota Leisure Time Physical Activity Questionnaire	“Would you say your mental health in general is excellent, very good, good, fair, or poor?” SF-8
Humphreys (2013)	Cross-sectional	United Kingdom	989 adults (68% female)	$R = 16+$	Leisure time, Work, Transport	Recent Physical Activity Questionnaire, 7 day travel to work recall	
Hyde (2011)	Cross-sectional	United States	190 young adults (66% female)	$M = 19.3 \pm 2.8$	Leisure time	The Godin Leisure-Time Exercise Questionnaire	20 items measuring four components of affect
Im (2014)	Cross-sectional	United States	542 middle aged adults (100% female)	$M = 49.04 \pm 6.05$	Leisure time, Work, Household	Kaiser Physical Activity Survey	Depression Index for Midlife Women
Inal (2007)	Cross-sectional	Instanbul	133 older aged adults (36% female)	$M = 73.9 \pm 8$	Leisure time	Items addressing frequency and duration	Life Satisfaction Index
Iwasaki (2001)	Cross-sectional	Canada	17,626 (sex not reported)	$R = 12+$ years	Leisure time	Physically active leisure index	Mental Distress Scale and Depression Scale
Jewett (2014)	Longitudinal	Canada	853 young adults (54% female)	$M = 20.4$	School sport	One item	One item for mental health, one item for stress, Major Depression Inventory
Joseph (2014)	Cross-sectional	United States	590 young adults (48% female)	$M = 20.4 \pm 1.7$	Leisure time	The Godin Leisure-Time Exercise Questionnaire	Satisfaction with Life Scale, Bradburn Affect Balance Scale
Jurakić (2010)	Cross-sectional	Croatia	989 adolescents and adults (51% female)	Not reported	Leisure time, Work, Transport, Household	IPAQ	SF-36
Kilpatrick (2013)	Cross-sectional	Australia	3,367 adults (72% female)	$M = 46.2$	Leisure time	IPAQ	Kessler Psychological Distress Scale
Kim (2014)	Cross-sectional	United States	1,708 middle and older aged adults (61% female)	$M = 67.88 \pm 11.17$	Leisure time	4 items related to the frequency of participation	5 items were borrowed from an existing study for life satisfaction, the degree of experiencing 7 descriptive adjectives. BDI-21
Korniloff (2012)	Cross-sectional	Finland	927 older aged adults (50% female)	$R = 65 - 74$	Leisure time	Frequency of physical activities in leisure-time at different ages.	
Kremer (2014)	Cross-sectional	Australia	8,256 adolescents (52% female)	$M = 11.5 \pm 0.8$	Leisure time, School sport, Physical education	Frequency of after school and weekend sport. The number of sports teams played on at school. Number of days per week attended physical education.	Short Mood and Feelings Questionnaire
Ku (2009)	Longitudinal	Taiwan	2,831 middle and older aged adults (46% female)	$R = 50 - 89$ (baseline)	Leisure time	“Do you usually engage in any type of leisure-time PA?”	10-item Chinese version of CES-D
Kull (2012)	Cross-sectional	Estonia	956 adults (100% female)	$R = 18 - 50$	Leisure time, Work, Transport	Global Physical Activity Questionnaire	BDI-21
Kwag (2011)	Cross-sectional	United States	163 older aged adults	$M = 81.84 \pm 7.05$	Household	Participation in lawn or yard work, gardening, or	Perceived Stress Scale

			(73% female)			heavy housework such as vacuuming.	Geriatric Depression Scale
Lee (2012)	Cross-sectional	United States	624 older aged adults (72% female)	M = 77.35	Leisure time, Household	Physical Activity Level	CES-D
Lin (2008)	Cross-sectional	United States	528 adults (46% female)	M = 45.4 ± 11.3	Leisure time, Work, , Household	3 day 24 hour recall	BDI-21
Lutz (2007)	Longitudinal	United States	203 adults (31% female)	M = 43.61 ± 9.79	Leisure time	Leisure-time Exercise Questionnaire	Percieved Stress Scale
Malabo (2007)	Cross-sectional	South Africa	293 adults (48% female)	M = 25.9 ± 4.20	Leisure time	The Physical Activity Index	Satisfaction with Life Scale, GHQ12, The Affectometer 2
Martins (2013)	Cross-sectional	Brazil	506 adults (100% male)	M = 29 ± 9.77	Leisure time, Transport	Baecke Questionnaire, Travel mode binary variable	Brazilian version of the GHQ-12
McKercher (2009)	Cross-sectional	Australia	1,995 young adults (52% female)	M = 31.4 ± 2.6 (females) M = 31.6 ± 2.6 (males)	Leisure time, Work	IPAQ	Composite International Diagnostic Interview
McKercher (2013)	Cross-sectional	Australia	1,995 adults (52% female)	M = 31.5 ± 2.6	Leisure time	IPAQ	DSM-IV Criteria for Major Depression
McKercher (2012)	Cross-sectional	Australia	6,070 adolescents (50% female)	R = 9 - 15	Transport, Physical education	Self-report duration and frequency of walking and cycling to and from school and physical education.	Negative Affect Scale of the Affect Balance Scale
Molarius (2009)	Cross-sectional	Sweden	34,645 adults (53% female)	R = 18 - 84	Leisure time	"How much do you exercise physically in your leisure time?"	One item: severity of anxiety or depression.
Moore (1999)	Cross-sectional	United States	146 middle and older aged adults (76% female)	M = 56.6 ± 6.6	Leisure time	Minnesota Leisure Time Physical Activity Questionnaire	BDI
Motl (2004)	Longitudinal	United States	4,594 adolescents (49% female)	M = 12.7 ± 0.4	Leisure time	"Do you get some regular physical activity outside of school?"	CES-D
Mouchacca (2013)	Longitudinal	Australia	1,382 adults (100% female)	M = 35.7 ± 7.7	Leisure time	IPAQ-Long Version	Perceived Stress Scale
Mouton (2000)	Cross-sectional	United States	210 older aged adults (61% female)	M = 72.12 ± 6.32	Leisure time	Minnesota Leisure Time Physical Activity Questionnaire	Geriatric Depression Scale
Mutrie (2007)	Cross-sectional	Scotland	1,742 adults (55% female)	Three age cohorts: approximately 24, 44, and 64	Leisure time, Work, Household	Amount and intensity of PA during leisure-time, work, and housework.	Hospital Anxiety and Depression Scale
Nakamura (2014)	Cross-sectional	Brazil	1,001 adults (58% female)	M = 46 ± 17 (females) M = 43 ± 16 (males)	Leisure time	IPAQ-Long Version	SF-36
Ohta (2007)	Cross-sectional	Japan	670 adults (36% female)	R = 18 - 60 years	Leisure time, Transport	State the weekly hours of leisure-time exercise engaged in. Frequency and duration of walking and bicycling.	GHQ12
Pagán (2014)	Longitudinal	Germany	53,819 adults (sex not reported)	R = 16+	Leisure time	"How often do you take part in active sports?"	"How satisfied are you with life, all things considered?"
Pasco(2011)	Cross-sectional	Australia	547 older aged adults (44% female)	R = 65 – 80	Leisure time	Subjects reported participation in habitual LT physical activity throughout the prior year.	The Structured Clinical Interview for DSM-IV Criteria for Major Depression Research

							Version, Non-patient edition
Pastor (2003)	Cross-sectional	Spain	1,038 adolescents (51% female)	$M = 16.31 \pm 0.92$	Leisure time	Frequency of sport participation not at school.	Two items: Frequency of feeling nervous and depressed.
Pedusic (2015)	Cross-sectional	Croatia	1,163 young adults (62% female)	$M = 21.5 \pm 1.8$	Leisure time, work, Transport, Household	IPAQ-Long Version	Satisfaction with Life Scale
Peeters (2014)	Cross-sectional	Australia	25,596 adults (100% female)	3 age cohorts: $M = 27.6 \pm 1.5$, $M = 52.2 \pm 1.5$, $M = 78.2 \pm 1.5$ $R = 11 - 15$	Leisure time, Household	Modified version of the Active Australia Questionnaire	SF-36
Petronyte (2009)	Cross-sectional	20 European countries	106,319 adolescents (51% female)		Leisure time	Amount of exercise during free time.	Health Behavior in School-aged Children symptom checklist.
Pickett (2012)	Cross-sectional	United Kingdom	164 adults (64% female)	$M = 30$	Leisure time	Leisure Time Exercise Questionnaire	BDI-II, PANAS
Purakom (2014)	Cross-sectional	Thailand	226 older aged adults (76% female)	$M = 68.41 \pm 6.42$	Leisure time, Work, Transport	Global Physical Activity Questionnaire	Mental Health Indicator short version 2007 (TMHI-15)
Rocha (2012)	Cross-sectional	Brazil	3,597 adults (71% female)	$R = 15+$	Leisure time	Frequency and intensity of PA during last month.	Self-Reporting Questionnaire (SRQ-20)
Sabiston (2013)	Longitudinal	Canada	860 adolescents (54% female)	$M = 12.7 \pm 0.5$ (baseline), $M = 20.4 \pm 0.7$ (last cycle)	Leisure time	The number of organized sports teams involved in during the last year.	"A validated depressive symptom scale"
Sanchez-Villegas (2008)	Longitudinal	Spain	10,381 adults (sex not reported)	Not reported	Leisure time	Leisure Time Physical Activity questionnaire collecting information about 17 activities.	Self-reported physician diagnosis of depression, anxiety, or stress, or the use of antidepressants or tranquilizers.
Sanchez-Villegas (2012)	Longitudinal	Spain	4,206 adults (sex not reported)	Not reported	Leisure time	Leisure Time Physical Activity questionnaire collecting information about 17 activities.	Spanish version of the SF-3
Sasidharan (2006)	Cross-sectional	United States	774 middle and older aged adults (sex not reported)	$R = 50+$	Leisure time	How many hours a day did you spend walking?	Single item from the Satisfaction with Life Scale, single item from the adapted Daily Hassles Scale
Schnohr (2005)	Cross-sectional	Denmark	12,028 adults (54% female)	$R = 20 - 79$	Leisure time	Frequency and intensity of PA.	Do you feel stressed? How often do you feel stressed? How satisfied have you been with your life within the last year?
Serrano-Sanchez (2013)	Cross-sectional	Spain	246 older aged adults (100% male)	$R = 65+$	Leisure time	Frequency and duration of PA and sport club affiliation.	10-item CES-D
Sieverdes (2012)	Cross-sectional	United States	9,580 adults (100% male)	$M = 48.9 \pm 9.7$	Leisure time	Type, frequency, and duration of leisure-time PA.	General Well-Being Schedule, CES-D, Bradburn's Affect Balance Scale, Health Opinion Survey
Stephens (1988)	Cross-sectional	Canada	23,791 adults (sex not reported)	$R = 15+$	Leisure time	Frequency and duration of leisure-time PA.	Two items from the Health Related Quality of Life Scale, a question
Surkan (2005)	Cross-sectional	United States	409 adults (59% female)	$R = 18+$	Leisure time	Non-work related physical activity in the past 30 days?	

							about satisfaction with life SF-36
Tessier (2005)	Longitudinal	France	3,891 adults (58% female)	M = 51.8 ± 6.1	Leisure time	Modifiable Activity Questionnaire	
Teychenne (2008)	Cross-sectional	Australia	1,501 adults (100% female)	M = 42 R = 18 - 65	Leisure time, Work, Transport, Household	IPAQ	30-item GHQ
Teychenne (2010)	Cross-sectional	Australia	3,645 adults (100% female)	R = 18 - 45	Leisure time, Work, Transport, Household	IPAQ	CES-D
Teychenne (2014)	Cross-sectional	Australia	4,065 adults (100% female)	R = 18 - 45	Leisure time	IPAQ-Long Version	10 item CES-D
Thøgersen-Ntoumani (2005)	Cross-sectional	United Kingdom	312 adults (35% female)	M = 34.11 ± 8.07	Leisure time	Baecke Habitual Physical Activity Questionnaire	Satisfaction with Life Scale
Valois (2004)	Cross-sectional	United States	4,758 adolescents (53% female)	Grade 9 - 12 Students	Leisure time, Transport, School sport	One item for leisure-time, The number of sports teams played on at school. Frequency of physical education, Self-report minutes spent exercising during physical education.	Students' Life Satisfaction Scale
Vuillemin (2007)	Cross-sectional	France	5,654 adults (59% female)	M = 47.6 ± 6.5 (females), M = 25.1 ± 4.6 (males)	Leisure time	Modifiable Activity Questionnaire	SF-36
Wang (2011)	Longitudinal	Canada	17,276 adolescents and adults (52% female)	R = 12+	Leisure time	Modified version of the Minnesota Leisure Time Physical Activity Questionnaire	Composite International Diagnostic Interview - Short Form
Wendel-Vos (2004)	Longitudinal	Netherlands	1,871 adults (53% female)	M = 43.3 ± 9.3 (females), M = 44.1 ± 9.2 (males)	Leisure time	Duration of leisure-time PA.	Health Related Quality of Life
Winjndaele (2007)	Cross-sectional	Belgium	2,616 adults (45% female)	M = 47.8 ± 12.3 (females), M = 49.3 ± 13.0 (males)	Leisure time, Transport, Household	Flemish Physical Activity Computerized Questionnaire	Perceived Stress Scale
Yang (2012)	Longitudinal	Finland	935 adults (51% female)	M = 38.4 ± 5.06 (females) M = 37.9 ± 4.97 (males)	Leisure time	Frequency and intensity of PA.	Modified version of the BDI
Zullig (2011)	Cross-sectional	United States	245 adolescents (55% female)	Year 7 and 18	Physical education	Frequency of physical education attendance.	One item

Note: For brevity only the first author is listed.

BDI, Beck Depression Inventory; CES-D, Center for Epidemiological Studies Depression Scale; GHQ, General Health Questionnaire; IPAQ, International Physical Activity Questionnaire; M, mean; Mo, Mode; PA, physical activity; PANAS, Positive and Negative Affect Scale; R, range; SF, Short Form Health Survey

Table S3.

Mental health variables measured within each PA domain.

Positive Mental Health					Mental ill-health					
	Mental Health	Mental Wellbeing	Life Satisfaction	Positive Affect	Mental ill-health	Stress	Life Dissatisfaction	Negative Affect	Depression	Anxiety
Leisure-time PA	Older Adults n = 3,798 (k = 3)	Adults n = 36,478 (k = 6)	Adults n = 7,695 (k = 9)	Adults n = 49,204 (k = 9)	Adults n = 151,278 (k = 13)	Adults n = 57,700 (k = 10)	Adults n = 12,028 (k = 1)	Adults n = 47,496 (k = 8)	Older Adults n = 10,033 (k = 9)	Adults n = 64,214 (k = 5)
	Adults n = 22,970 (k = 8)		Adolescents n = 62,981 (k = 3)		Adolescents n = 108,989 (k = 3)				Adults n = 141,175 (k = 26)	Adolescents n = 3,050 (k = 2)
	Adolescents n = 8,726 (k = 2)				Children n = 13,476 (k = 1)				Adolescents n = 17,953 (k = 7)	
Work-related PA	Older Adults n = 226 (k = 1)	Adults n = 3,183 (k = 2)	Adults n = 1,750 (k = 1)		Adults n = 506 (k = 1)				Adults n = 47,910 (k = 7)	Adults n = 38,743 (k = 1)
	Adults n = 2,055 (k = 2)								Adolescents n = 1,742 (k = 1)	
Transport PA	Older Adults n = 226 (k = 1)	Adults n = 21,168 (k = 3)	Adults n = 1,750 (k = 1)		Adults n = 1,919 (k = 1)	Adults n = 4,535 (k = 2)		Adolescents n = 6,070 (k = 1)	Adults n = 6,102 (k = 3)	
	Adults n = 2,055 (k = 2)		Adolescents n = 4,758 (k = 1)		Adolescents n = 670 (k = 1)					
Household PA	Adults n = 2,055 (k = 2)	Adults n = 27,793 (k = 2)	Adults n = 1,750 (k = 1)		Adults n = 21,761 (k = 2)	Older Adults n = 163 (k = 1)			Older Adults n = 961 (k = 3)	
						Adults n = 4,535 (k = 2)			Adults n = 6,216 (k = 4)	
School Sport	Adolescents n = 853 (k = 1)		Adolescents n = 4,758 (k = 1)			Adolescents n = 853 (k = 1)			Adolescents n = 1,742 (k = 1)	
									Adolescents n = 8,700 (k = 2)	
Physical Education			Adolescents n = 5,003 (k = 2)					Adolescents n = 6,070 (k = 1)	Adolescents n = 7,847 (k = 1)	

Note: Blank cells indicate that no study examined the relationship between that particular physical activity domain and mental health variable. PA, physical activity

1 Table S4.

2 *Risk of bias of included studies.*

Author	Random selection of schools or participants	Eligibility criteria stated and study sample described	Valid assessment of physical activity	Valid assessment of mental health or mental ill-health	Covariates adjusted for in analyses	Power calculation reported and study adequately powered.	Risk of Bias Score /6
Aldana (1996)	0	1	0	1	1	0	3
Armstrong (2004)	1	0	1	1	0	1	4
Asztalos (2009)	1	1	1	1	1	0	5
Balboa-Castillo (2011)	1	1	1	1	1	0	5
Barrington (2012)	1	1	0	1	1	0	4
Bernaards (2006)	0	0	0	1	0	0	1
Bertheussen (2011)	1	1	1	1	1	0	5
Binsinger (2006)	0	1	0	1	0	0	2
Birkeland (2009)	0	1	1	1	1	0	4
Bogaert (2014)	1	1	1	1	1	0	5
Borges (2013)	0	0	1	1	1	0	3
Branco (2014)	1	1	1	1	1	0	5
Brooks (2014)	1	0	1	0	0	0	2
Brunes (2013)	0	1	1	1	1	0	4
Brunes (2014)	0	1	1	1	1	0	4
Bültmann (2002)	0	1	0	1	1	0	3
Bustamante (2013)	0	1	1	1	0	0	3
Buttery (2014)	1	1	0	1	0	0	3
Cerin (2009)	1	1	1	1	1	0	5
Chen (2012)	0	1	1	1	1	0	4
Chi (2015)	0	1	0	1	0	0	2
Choi (2008)	0	1	1	1	1	0	4
Craike (2010)	1	1	0	1	1	0	4
Desha (2007)	0	1	0	1	1	0	3
Doerksen (2014)	0	0	0	1	0	0	1
Dupuis (1995)	0	1	0	1	0	0	2
Eime (2014)	0	1	0	1	1	1	4
Feuerhahn (2014)	0	1	0	1	1	0	3
Fisher (2004)	1	1	1	1	0	1	5
Fløtnes (2011)	0	1	1	1	1	0	4
Giacobbi Jr (2007)	0	1	1	1	0	0	3
Goldfield (2011)	0	1	1	1	1	0	4
Griffiths (2010)	0	1	0	1	1	0	3

Hamer (2009)	0	1	1	1	1	0	4
Hamer (2009)	1	1	1	1	1	0	5
Herman (2015)	0	1	0	0	1	0	2
Humphreys (2013)	0	1	1	1	1	0	4
Hyde (2011)	0	1	1	1	0	0	3
Im (2014)	0	1	1	1	0	1	4
Inal (2007)	0	1	0	1	0	0	2
Iwasaki (2001)	0	0	0	0	1	0	1
Jewett (2014)	0	1	0	1	1	0	3
Joseph (2014)	0	1	1	1	0	0	3
Jurakić (2010)	1	0	1	1	1	0	4
Kilpatrick (2013)	0	1	1	1	1	0	4
Kim (2014)	0	1	0	0	0	0	1
Korniloff (2012)	1	1	0	1	1	0	4
Kremer (2014)	1	1	1	1	1	0	5
Ku (2009)	1	1	0	1	1	0	4
Kull (2012)	1	1	1	1	0	0	4
Kwag (2011)	0	1	1	1	0	0	3
Lee (2012)	0	1	0	1	1	0	3
Lin (2008)	0	1	1	1	1	0	4
Lutz (2007)	0	1	1	1	1	0	4
Malabo (2007)	0	1	0	1	0	0	2
Martin (2014)	0	1	0	1	1	0	3
Martins (2013)	0	1	1	1	1	0	4
McKercher (2009)	0	1	1	1	1	0	4
McKercher (2013)	1	1	1	1	1	0	5
McKercher (2012)	1	1	0	1	1	0	4
Molarius (2009)	1	1	0	0	1	0	3
Moore (1999)	0	1	1	1	1	0	4
Motl (2004)	1	1	1	1	1	0	5
Mouchacca (2013)	1	1	1	1	1	0	5
Mouton (2000)	0	1	1	1	0	0	3
Mutrie (2007)	1	1	0	1	1	0	4
Nakamura (2014)	1	1	1	1	1	0	5
Ohta (2007)	0	1	1	1	1	0	4
Pagán (2014)	0	0	0	0	0	0	0
Pasco (2011)	1	1	1	1	0	0	4
Pastor (2003)	1	1	0	0	0	0	2
Pedisic (2015)	1	1	1	1	1	1	6
Peeters (2014)	1	1	1	1	1	1	6
Petronyte (2009)	0	1	0	1	1	0	3
Pickett (2012)	0	1	0	1	1	0	3
Purakom (2014)	0	1	1	1	1	0	4

Rocha (2012)	0	1	0	1	1	1	4
Sabiston (2013)	0	1	0	1	1	0	3
Sanchez-Villegas (2008)	0	0	1	0	1	0	2
Sanchez-Villegas (2012)	0	0	1	1	1	0	3
Sasidharan (2006)	0	0	0	0	0	0	0
Schnohr (2005)	1	1	0	0	1	0	3
Serrano-Sanchez (2013)	1	1	0	1	0	0	3
Sieverdes (2012)	0	1	1	1	1	0	4
Stephens (1988)	0	0	0	1	1	0	2
Surkan (2005)	0	1	0	0	1	0	2
Tessier (2005)	0	1	1	1	1	1	5
Teychenne (2008)	1	1	1	1	1	0	5
Teychenne (2010)	1	1	1	1	1	0	5
Teychenne (2014)	1	1	1	1	1	0	5
Thøgersen-Ntoumani (2005)	0	1	1	1	0	0	3
Valois (2004)	1	0	0	1	1	0	3
Vuillemin (2007)	1	1	0	1	1	0	4
Wang (2011)	1	1	0	1	1	0	4
Wendel-Vos (2004)	1	1	1	1	1	0	5
Winjndaele (2007)	1	1	1	1	0	0	4
Yang (2012)	1	1	1	1	1	0	5
Zullig (2011)	0	1	1	0	1	0	3

3 *Note:* For brevity only the first author is listed.

4 1, present and explicitly explained; 0, absent or inadequately discussed

5

6

7

8

Mental ill-health **Mental health**

a) Leisure-time Physical Activity

g) Leisure-time Physical Activity

b) Work-related Physical Activity

h) Work-related Physical Activity

c) Transport Physical Activity

i) Transport Physical Activity

d) Household Physical Activity

j) Household Physical Activity

Figure S1. Individual study effect size estimates.